

Ecosystem Services and Southern Wetland Forests:

Baseline Value and Future Prospects

September 2017

Spencer Phillips, Ph.D.

John Stoner

JP Schmidt, Ph.D.

Sam Davis, Ph.D.

Prepared for:

Dogwood Alliance



KEY-LOG.
economics_{LLC}

Research and strategy for the land community.

Keylogeconomics.com

Executive Summary

Some 35 million acres of wetland forests provide valuable ecosystem services for people living in the U.S. South¹ and beyond. These services include water filtration, carbon sequestration and local climate regulation, waste assimilation, protection from extreme events, and recreational opportunities that, when measured in monetary terms, represent more than \$503.8 billion per year in market and non-market value. How much of this value will be available in the future hinges on how much of this forest remains and how it is managed.

We consider the allocation those 35 million acres of wetland forest to various land use/land management categories in two future scenarios and estimate the ecosystem service value in each, in addition to the 2016 baseline. In both the “business-as-usual” (“BAU”) and the “Conservation” scenario, some wetland forest acreage will be converted to non-forested uses, some will be placed in protective designations and managed primarily for biodiversity and other non-commodity values, while other areas will remain available for timber harvest. Among these latter areas, some land will be in more intensive forest management, including plantations, some will be managed under “ecological forestry” standards, and some may continue in a state of benign neglect. Overlapping all areas except those converted to non-forest uses, are streamside-management zones with widths as currently defined in state-by-state regulations for the baseline and BAU scenario, but expanded to 150’ in the conservation scenario.

Wetland forests in each of the land use/land management categories will produce an array of ecosystem services, but not all at the same rate per acre. Land converted to urban/developed uses, for example, will provide less in the way of flood damage protection or recreational opportunities than the natural forests they replace. And wetland forest land allocated to more intensive forest management will produce more timber, but perhaps less carbon storage, water supply, or protection from extreme events, like flooding.

To the extent possible, our scenarios incorporate spatially explicit information on likelihood of conversion, proposals for expansion of protected areas, and the geographic reach of forest management regulations (specifically streamside management zones). Other facets of the baseline conditions and future scenarios, however, can only be applied to estimates of residual acreage—that is, acreage not geographically tied to any particular cause or type of land use/forest management change. Our procedures and techniques, primarily spatial analysis using ArcGIS², for allocating wetland forests to land use/land management categories are described in detail in the [Spatial Analysis Methods](#) section below.

With acreage estimates in hand, we proceed to estimate the economic value of these forests today and in the plausible future scenarios, we apply the “benefits transfer method” (“BTM”) to the areas identified in each scenario. According to this method, acreage is multiplied by per-acre ecosystem service values gleaned from existing literature, including the TEEB (“The Economics of Ecosystems and Biodiversity”) database (Van der Ploeg, Wang, Gebre Weldmichael, & De Groot, 2010). These source studies typically provide a range of value-per-acre estimates for various sets or arrays of ecosystem services for one or more land cover type(s).

Unfortunately little literature exists regarding the differential ecosystem service productivity—that is, the per-acre values—for different management regimes within a land cover type. It is difficult to speculate on just how much more (or less) water supply, timber, or recreational value would be provided to different management

¹ For purposes of this study, this region comprises Virginia, North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, Texas, Oklahoma, Arkansas, Missouri, Kentucky, and Tennessee. See maps, below.

² We are grateful to ESRI and its [Conservation GIS Program](#) for generous support in the form of the ArcMap used in this analysis.

types within the forest, and especially, the wetland forest land use/land cover category. We can, however, speculate with some confidence regarding the relative productivity of different management types. Land managed for conservation purposes, where commercial resource extraction including timber harvest is prohibited, would produce no raw material (timber) value, whereas land in the “ecological forestry” category and “intensive forestry” would provide more raw materials.

To overcome this limitation in the available data while still developing estimates of ecosystem service value, we have used the available ranges of ecosystem service productivity estimates and assigned different values from within those ranges to our land management types on an ecosystem service-by-ecosystem service basis. We describe the details of these assignments under [Ecosystem Service Valuation](#) in the Methods section, below.

The internal logic of this method, which we believe corresponds to likely real-world outcomes, is that differences among the scenarios in terms of total ecosystem service value are driven by differences in the number of acres allocated to different land-use/land-management categories, and by differences in ecosystem services productivity among the land-use/land-management categories.

In brief, we estimate a fairly low level of conversion of wetland forests to non-forest uses and, due to potential implementation of land protection priorities, an increase in overall ecosystem service value in both the business-as-usual and conservation scenarios. Ecosystem service value is further boosted in the conservation scenario due to an expansion of streamside management zones.

Conversion of wetland forests to urban open space and urban developed areas would mean a loss of 407,334 wetland forest acres in the business-as-usual scenario and a loss of 362,937 acres in the conservation scenario. Additional acreage in each scenario would be converted from wetland forest to pine plantation, leaving 34.4 and 34.5 million acres as wetland forests in the BAU and conservation scenarios, respectively. (See Table 1.)

Table 1. Summary of Land-Use/Land Management Elements of Baseline, Business-as-Usual, and Conservation Scenarios, with Acreage and Estimated Ecosystem Service Value

Land Use / Land Management Category	Allocated Wetland Forest Acreage and Ecosystem Service Value (2016\$)		
	Baseline	Business as Usual	Conservation
Totals^a:	35.1 million acres (all wetland forest) ESV: \$503.8 billion	35.1 million acres (34.4 mm acres are wetland forest) ESV: \$526.9 billion	35.1 million acres (34.5 mm acres are wetland forest) ESV: \$549.1 billion
Protected Areas	Protected Areas Database: GAP category 1 and 2 lands, which are permanently protected from development and managed for values other than commodities. Vegetation management for the purposes of maintaining or enhancing other resource values is permitted. Commodity harvest is not permitted.		
	Baseline Protection: 2,812,908 acres Total Ecosystem Service Value: \$51.1 billion		
Added to Protected Areas	n/a	Southeast Conservation Adaptation Strategy ("SECAS") Blueprint: 50% of "high" priority areas	SECAS Blueprint: 100% of "high" areas
		6,476,416 acres \$118.1 billion	12,952,832 acres \$236.3 billion
Converted to non-forest use	n/a	Areas at >50% risk of urbanization by 2050 (SLEUTH projections)	
		407,334 acres \$1.4 billion	362,937 acres \$1.3 billion
Streamside Management Zones ("SMZs")	Width varies by state ³ . Harvest may be permitted within SMZs.		150' for permanent/intermittent streams. No harvest in SMZs.
	70,874 acres w/in protected areas; 1,740,732 acres outside	377,953 acres w/in protected areas 1,415,374 acres outside	2,216,129 acres w/in protected areas 3,704,284 acres outside

³ For some states, the recommended SMZ width is a function of the slope. We are not including that variation in our model.

	protected areas ESV outside protected areas: \$31.6 billion	protected areas ESV outside protected areas: \$25.7 billion	protected areas ESV outside protected areas: \$67.3 billion
--	--	--	--

Table 1 continues on the next page.

Table 1, continued.

Land Use / Land Management Category	Allocate Wetland Forest Acreage		
	Baseline	Business as Usual	Conservation
Ecological Forestry (outside SMZ) 18633\$ / ACRE	Areas under management that meets certification standards. Scenario-specific SMZ in effect. Expect less timber value, more of other ecosystem service values.		
	453,776 acres ESV: \$8.3 billion	890,188 acres ESV: \$16.2 billion	1,032,274 acres ESV: \$18.8 billion
Intensive Forestry (outside SMZ) 1194\$ / ACRE	Forestry emphasizes fiber production. Scenario-specific SMZ in effect. No other distinguishing restrictions. Expect more timber value, less of other ecosystem service values.		
	7,826,391 acres ESV: \$9.1 billion	5,845,378 acres ESV: \$6.8 billion	4,667,465 acres ESV: \$5.5 billion
Plantation Forestry (outside SMZ) 1194\$ / ACRE	Wetland forests that have been drained and planted in (typically) pine and managed for fiber commodities. Scenario-specific SMZ in effect. Expect more timber value, less of other ecosystem service values.		
	0 acres, ESV: \$0	307,651 acres ESV: \$359.6 million	245,646 acres \$287.1 million
Benign Neglect 18550\$ / ACRE	Residual Category for acreage left after other allocations are made.		
	22,235,017 acres ESV: \$403.7 billion	16,913,573 acres ESV: \$307.1 billion	9,290,466 acres ESV: \$168.7 billion

Note:

- Because we track the disposition and ecosystem service production from all acres in our baseline scenario, the total acreage in all land use/land management categories stays constant. The lower wetland forest

acreage estimates (in parentheses on the “Totals” line for the BAU and conservation scenarios) are total acreage less acreage converted to urban uses and acreage converted to plantation forestry.

Collectively, these 35.1 million acres generate or support \$503.8 billion per year in diverse ecosystem services. Due to increases in conservation land, on which productivity gains for non commodity ecosystem services more than offset the loss of timber value, annual ecosystem service delivery increases to \$526.9 billion per year in the BAU scenario and \$549.1 billion per year in the conservation scenario. In each scenario, approximately 20% of total ecosystem service value is due to aesthetic value. Protection from extreme events, water flow regulation, and food supply about 15% each, and water supply and waste treatment contribute up 11.7 and 11.2 percent, respectively, of the total. The balance of ecosystem service value is delivered as air purification, climate regulation, erosion control, pollination, raw materials (timber), recreation, and soil formation. (See Table 2.)

Table 2. Annual Ecosystem Service Value in Baseline, Business-as-Usual, and Conservation Scenarios, in millions of 2016 dollars.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	503,850.0	526,873.0	549,121.6
Aesthetic Value	101,893.1	106,982.9	111,729.7
Air Quality	4,262.2	4,332.5	4,407.2
Climate Regulation	12,438.8	13,338.4	13,888.2
Erosion Control	1,947.1	2,031.2	2,118.0
Protection from Extreme Events	77,151.4	80,623.6	84,068.8
Food	70,682.5	73,955.5	77,278.3
Pollination	2,079.1	2,172.8	2,268.5
Raw Materials	1,370.8	1,652.9	1,976.5
Recreation	24,135.4	25,689.9	26,778.2
Soil Formation	18,794.9	18,576.5	18,600.3
Waste Treatment	56,215.5	58,810.1	61,446.2
Water Supply	59,135.1	61,981.6	64,738.4
Water Flow Regulation	73,744.0	76,725.2	79,823.4

Differences between the baseline and future scenarios arise due to differences among the number of acres in each land use/land management category and assumptions about the relative ecosystem service productivity of each land use category.

Contents

Executive Summary	1
Contents	7
Acronyms and Definitions	9
About the Authors	10
Study Objectives	11
Scenarios	11
Baseline	12
Business-as-Usual:	12
Conservation:	13
Methods	13
Methods for Land Allocation	14
Wetland Forests	14
Protected Areas	15
Urban Development	16
Streamside Management Zones	17
Canopy Cover	18
Contextual Information	20
Average Tree Age	20
Carbon Storage	20
Methods for Ecosystem Service Valuation	22
Ecosystem Service Estimates	26
Conclusions	28
Works Cited	28
APPENDIX 1: State-by-State Estimates	35
Alabama	36
Arkansas	38
	7

Florida	40
Georgia	42
Kentucky	44
Louisiana	46
Mississippi	48
Missouri	51
North Carolina	54
Oklahoma	56
South Carolina	58
Tennessee	60
Texas	62
Virginia	64

Appendix 2: Candidate Ecosystem Service Productivity Values for Land-Use and Ecosystem Service Combinations 66

Acronyms and Definitions

Baseline Scenario: The extent of wetland forests and their allocation among land use/land management categories, based on the most recent spatial data. (See [Baseline](#), page 11, for details.)

Business-as-Usual Scenario: The projected allocation of wetland forests (as identified in the Baseline) among various land use/land management categories in the year 2050 if current trends continue and one half of the acreage identified as “high priority” for conservation is placed into a protective status. (See [Business-as-Usual](#), page 12, for details).

BTM: Benefit Transfer Method, a method for estimating the value of ecosystem services in a study region based on values estimated for similar resources in other places

Conservation Scenario: The projected allocation of wetland forests (as identified in the Baseline) among various land use/land management categories in the year 2050 with greater conservation efforts than those in the Baseline and Business-as-Usual Scenarios. (See [Conservation](#), page 12 for details)

Ecological Forestry: Areas under management that meet certification standards with an emphasis on supporting and protecting native ecosystem health and production.

Ecosystem Services: Benefits people derive from ecosystems. Specifically, and as modeled in this report, they comprise the monetary value of various benefits that, were they not provided by nature, people would have to provide for themselves. Please see table 5 for a list of the specific services included in this study.

GAP 1&2: U.S. Geological Survey’s Gap Analysis Program, provides information and designations for areas that are currently protected for biodiversity.

Protected Status: Areas which are permanently protected from development and managed for values other than commodities which includes GAP 1&2 areas as well as scenario specific portions of the SECAS protection designation (100% of “High” SECAS priority for the Conservation Scenario, and 50% of the “High” SECAS priority for the BAU Scenario).

SECAS: Southeast Conservation Adaptation Strategy, a multi-state, multi-agency effort to create and share restoration and protection designations for the Southeast United States and the Caribbean.

SMZ: Streamside Management Zones consist of land on either side of permanent or intermittent streams designated by the U.S. Geological Survey. Each scenario provides different widths and land uses permitted in the SMZ (state specific widths in the BAU and Baseline scenarios, and 150 feet in the Conservation Scenario) (U.S. Geological Survey, 2017).

About the Authors

Spencer Phillips, PhD has been conducting and directing applied research into the relationships between natural resource stewardship, environmental quality, and human well-being for more than 25 years. He was a staff economist first at the White House Council on Environmental Quality during the first Bush Administration, and then at The Wilderness Society, where he later served as Vice President for Ecology and Economics Research. Dr. Phillips founded Key-Log Economics to help ensure that sound, independent economic research is available to those working to solve the critical environmental problems of our time. He is also a lecturer in economics, natural resource policy, and GIS analysis at the undergraduate and graduate level. Phillips holds a B.A. in economics from the University of Virginia and an M.S. and Ph.D. from Virginia Tech.

Nick Stoner is a masters candidate in the Yale School of Forestry & Environmental Studies focusing on business and the environment. He also holds a B.S. in biology from James Madison University and has field and laboratory research experience for NOAA and other organizations.

JP Schmidt, PhD is a Research Scientist at the Odum School of Ecology at the University of Georgia, where he also received his doctorate. JP's research background lies in vegetation monitoring, land use change, and plant population biology. More recently, however, his research focus has expanded to include the ecology of infectious disease and work with economists on the valuation of ecosystem services.

Sam Davis, PhD has been the Research and Program Manager at Dogwood Alliance since early 2016. Sam earned a Ph.D. in Environmental Science in 2015 at Wright State University and completed a postdoc at University of California Merced. Sam's previous research areas include plant-insect interactions, exotic species impacts, forest gap modeling, and environmental justice issues.

Key-Log Economics remains solely responsible for the content of this report, the underlying research methods, and the conclusions drawn. We have used the best available data and employed appropriate and feasible estimation methods but nevertheless make no claim regarding the extent to which these estimates will match the actual magnitude of economic effects experienced under future scenarios for regional wetland forest conservation and management.

Study Objectives

The wetland forests of the southern United States support biodiversity, sequester and store carbon, filter drinking water, protect people and property from losses in floods and storms, and otherwise provide value to people. Over the course of centuries, millions of acres of these forests have been drained and converted for use as cropland, for plantation forestry, and for housing and other developed uses (Dahl, 2011). The pace of conversion has slowed in recent decades as land protection efforts have prioritized areas with higher ecological values and as water-quality-based restrictions on wetland conversion have come into effect. Today, wetland forest protection may be even more important as a means of ensuring the continued provision of ecosystem services that will be needed more as climate change increases the need for services wetland forests are well suited to provide, such as protection against extreme weather events, and as places to sequester and store carbon or to preserve “critical natural capital” (Farley, 2012) that might otherwise be lost regionally or globally.

Information about the potential economic benefits of wetland forest conservation can help inform decisions about whether, where, and by what means to pursue that conservation. Our objective is to provide initial estimates of those benefits and a starting framework for future analysis as new information becomes available that can also be “down-scaled” to assess ecosystem service value at a sub-regional, state, or even project level using more detailed information about specific forest composition and other local conditions.

Our method, has three primary steps. First, we use GIS analysis and other data to identify the current extent and location of wetland forests across the U.S. South and to estimate how these areas are allocated among several land use/land management categories. This step provides our “baseline scenario” to for comparison to two possible future scenarios. Second, we develop those two future scenarios—labeled “business-as-usual” and “conservation”—and estimate the allocation of wetland forest acreage to the same array of land use/land management categories. (We selected 2050 as a timeframe for these scenarios in part for the availability of projections regarding potential conversion to non-forested uses and because the 30+ years between now and then may provide sufficient time to attain other conservation objectives embodied in the scenarios.) Third, we apply the benefits-transfer method (“BTM”) to develop estimates of the annual value of 13 ecosystem services delivered by the areas identified as wetland forests in the baseline. Differences in ecosystem service value between the baseline and future scenarios arise due to projected differences in the number of acres allocated to the various land use/land management categories as well as too differences among the land-use/land management categories in terms of per-acre ecosystem-service productivity.

Scenarios

The question of how much ecosystem service value wetland forests provide, or could provide in the future, depends on the extent of these forests and the use or management to which they are subject. is tied to questions of how extensive these forests are now, how they are used now, and how that extent and use may change in the future. We define the extent of wetland forests and their allocation to land use/land management categories for three scenarios: Baseline (i.e., current conditions); Business-As-Usual, or BAU; and Conservation.

Please refer to Table 1, above, which lays these scenarios out schematically and provides the estimated acreage and and ecosystem service value for each scenario and in each land-use/land-management category. Details on the spatial analysis conducted to evaluate each scenarios, along with sample maps, are provided under [Spatial Analysis Methods](#), below

Baseline

Following the method outlined by the Natural Resources Defense Council (2015), we use satellite data to identify the current extent and location of wetland forests. Specifically, wetland forests are defined here as those areas identified as “woody wetlands” in the National Land Cover Database that are also identified both as as “forests” in data from the U.S. GAP program and as “tree-dominated” in the USDA Forest Service’s LANDFIRE program (Fry, et al., 2011; National Gap Analysis Program, 2015; and LANDFIRE Program, 2017). Some 35.1 million acres meet these criteria.

Further GIS analysis using additional information from the GAP program, specifically the Protected Areas Database, identifies the extent to which wetland forest acreage is already protected. “Protected” is defined for our purposes as being classified as an area in GAP status 1 or 2. As defined by the program,

Status 1 [areas have] permanent protection from conversion of natural land cover and a mandated management plan in operation to maintain a natural state within which disturbance events (of natural type, frequency, intensity, and legacy) are allowed to proceed without interference or are mimicked through management.

Status 2 [areas have] permanent protection from conversion of natural land cover and a mandated management plan in operation to maintain a primarily natural state, but which may receive uses or management practices that degrade the quality of existing natural communities, including suppression of natural disturbance.

In contrast to GAP Status 3 lands, these Status 1 and 2 lands are not “subject to extractive uses of either a broad, low-intensity type (e.g., logging, OHV recreation) or localized intense type (e.g., mining) (National Gap Analysis Program, 2015, metadata).”

Streamside Management Zones are classified as areas surrounding permanent and intermediate streams designated by the U.S. Geological Survey (U.S. Geological Survey, 2017). SMZ width in the Baseline Scenario corresponds with the state specific SMZs that are in effect today, and the ability to harvest materials out of these areas remains in place. Wetland forests that fall within these SMZ’s are subject to harvesting as long as they do not fall within the designated protected areas of GAP Status 1 and 2.

To define areas designated as ecological forestry and benign neglect, canopy cover was used as a defining factor. We identified these areas as those with 66% or higher canopy cover. Certified acreage was collected from the Forest Stewardship Council and we found that ~2% of region’s forests are under certified forestry of one sort or another (Forest Stewardship Council, 2016). We defined Ecological forestry acreage within the Baseline Scenario as 2% of unprotected area, which also had 66% canopy coverage. Benign neglect is the total unprotected wetland forest acreage with 66% canopy closure, less the ecological forestry portion. Intensive forestry, where areas of wetland forest are more likely to be managed intensively for timber, are defined as areas with less than 66% canopy coverage and that fall outside of protected GAP 1 and 2 areas.

Business-as-Usual:

The BAU Scenario projects where wetland forests could be (from a land-use/land-management perspective) in 2050 in our study region. This projection is relative to the Baseline Scenario.

Some areas in the BAU scenario are likely to be converted to non-forest (urban) uses by the year 2050. Acreage within these areas are allocated to “urban developed” and “urban open space”, which are the subcategories for

which distinct ecosystem service values are available, according to the current distribution of those sub-categories across the region (Belyea & Terando, 2014).

Protected areas increase in the BAU scenario. We assume that 50% of the high priority areas identified by SECAS will be added to the region's GAP 1 and 2 lands by 2050. We foresee no changes in the size or accessibility to harvesting of the SMZs in the BAU Scenario. We assume an increase in the Ecological forestry percentage to 5%, gaining new area from the benign neglect category. We further assume that 5% of the remaining wetland forest areas (i.e., areas that would otherwise be intensively managed for timber) are converted to pine plantations, effectively leaving the wetland forest type.

Conservation:

This Scenario also defines where wetland forests could be in 2050 relative to the baseline. The Conservation Scenario entails greater protection and more careful stewardship of wetland forests compared to the BAU Scenario.

The areas that are likely to be converted to non-forest uses by 2050 remain the same in the Conservation Scenario, but the protected lands are increased to include all high-priority lands identified in SECAS as well as the currently protected (GAP 1 and 2) lands.

In the Conservation Scenario SMZs are increased to 150 feet on both sides of permanent and intermittent streams in every state, and SMZs are, effectively, a no-harvest zone. We also assume that 10% of the region's wetland forests are, by 2050, under ecological forestry in the Conservation Scenario, reducing the number of acres falling under the more intensive management categories as well as benign neglect. As in the BAU, scenario, we assume that 5% of the land outside protected areas, not in an SMZ, not converted to urban uses, and not in ecological forestry or benign neglect would be put into pine plantations.

Methods

Estimation of ecosystem service value requires two general steps:

1. Allocate total wetland forest acreages to one of eight land-use/land-management categories for which one would expect differing productivity for different ecosystem services. We perform this allocation for the baseline and the two future scenarios, BAU and Conservation.
2. Multiply acreage in each land-use/land-management category by the ecosystem service value per acre for each of [13] individual ecosystem services. For those ecosystem services which have, based on literature review, higher or lower productivity would be expected, we have applied land use-specific factors to the acreage each land use. For example, raw materials (i.e., timber), and climate regulation are two services for which productivity would be expected to differ between conservation areas and areas managed intensively for timber.

The result is a three-dimensional dataset with dollar-value estimates of ecosystem services for 13 ecosystem services harbored or produced within 8 land uses under the 3 scenarios (including the baseline). These atomized value estimates can then be summed in any dimension and compared to illuminate, for example, how total climate regulation value varies between the BAU and Conservation scenarios. In Appendix 1, we present state-by-state estimates of ecosystem service value of wetland forests.


Methods for Land Allocation

This project relied heavily on GIS analysis to identify and to calculate the area associated with each of the land use/land management scenarios (Baseline, Business as Usual, and Conservation). Six separate land use allocations (Protected Area, Urban Development, Streamside Management Zones, Tree Age, Canopy Cover, and Carbon Sequestration) were calculated for each of the scenarios and used to estimate the area of each land use type and/or to estimate key attributes used to allocate residual acreage—the area of wetland forest not tied to specific places on the map—among different forest management regimes.

Wetland Forests

The first step of the data analytics for this study entailed designating which areas should be considered wetland forest within the study region. Without knowing which areas were wetland forests, no ecosystem service data could be connected to those forests. By using a combination of the National Land Cover Database (U.S. Geological Survey, 2014), the National Gap Analysis Program (National Gap Analysis Program, 2015), and the LANDFIRE program (LANDFIRE Program, 2017), wetland forests were successfully defined for our study region (Figure 1). Once a wetland forest map layer was produced, it could be used to define all of the other land use allocations. This layer is the foundation for the ecosystem service values, and the bedrock of this study.


Figure 1: Wetland forests and in the U.S. South



Protected Areas

Under each of the conservation scenarios, protected areas consist of both GAP 1 and 2 lands, and Southeast Conservation Adaptation Strategy (“SECAS”) blueprint areas. GAP 1 and 2 lands are permanently protected from development and managed for values other than commodities. SECAS areas are designated as having either “high” or “medium” conservation value. For the BAU scenario, the GAP 1 and 2 lands as well as 50% of the SECAS high areas are considered protected going forward. In the Conservation scenario, the GAP 1 and 2 lands and 100% of the SECAS High areas are considered protected. For this report the SECAS “medium” designation (outside of GAP 1 and 2 lands) is added for context but does not fall within either conservation scenario’s protection.

GIS data layers were obtained for both GAP 1 and 2 (National Gap Analysis Program, 2015), and SECAS (Snider, 2016). Each layer was altered to fit the study region and then combined to indicate all the areas that have protection from GAP 1 and 2, SECAS, or both. Once the protection areas were defined, the wetland forest data from the National Land Cover Database (Fry et al, 2011) was added to produce the area of wetland forest under each protection designation (Figure 2).

Figure 2: Wetland forest within protected areas

Source: U.S. Geological Survey, 2014; National Gap Analysis Program, 2015; Snider, 2016; LANDFIRE Program, 2017


Urban Development

One thing that is almost a certainty when producing future scenarios is human expansion. Urban development is crucial to understand when creating the BAU and Conservation scenarios. To account for this future development a SLEUTH (Slope, Land use, Excluded, Urban, Transportation and Hillshade) database was aligned with the study region to predict areas that had a probability of 50% or greater of being developed by 2050 (Belyea & Terando, 2014) (Figure 3).

The urban development data layer was then cross-referenced with wetland forest data as well as streamside management zones, and protected areas. This produced areas of wetland forest, and streamside management zones that will be developed under each scenario. The assumption was made that unless the development falls within one of the designated protected areas created above, it will be converted into the urban land use designation and no longer supply ecosystem services. It should also be known that development areas that fall within streamside management zones were regarded as developed due to the fact that they are not protected at the time of this study. The urban land use designation contains 2 subcategories: “urban open space” and “urban developed”. These subcategories were defined by the ratio of existing urban open space, and urban developed (U.S. Geological Survey, 2014).

Figure 3: Urban areas and potential urbanization before 2050

Areas with 50% or greater probability of conversion to urban uses by 2050, including areas identified as wetland forests.




Streamside Management Zones

One of the largest differences between the BAU and Conservation scenarios is the size of streamside management zones (SMZ). Under the BAU scenario, the SMZs are state specific on either side of a stream, and harvesting is permitted within these zones. Under the Conservation scenario the SMZs span 150 feet on either side of a stream and no harvesting is permitted.

To accomplish this, stream data was acquired from national hydrography data (U.S. Geological Survey, 2017) and buffers were created around the streams depending on the buffer size under each scenario (state specific (BAU) and 150 feet (Conservation)) (Figure 4). Once the SMZs were defined, the area within them was combined with

the existing protection, wetland forest, and urbanization layers to find the area of each designation within the SMZ. This was done for both the state specific SMZs and the 150 Foot SMZs.

Figure 4: Streamside management zones, both BAU (State Specific) and Conservation (150 feet) scenarios
SMZ interactions with wetland forests within each conservation scenario, as well as the wetland forests outside of the SMZs to provide context.



Sources: U.S. Geological Survey, 2014; National Gap Analysis Program, 2015; LANDFIRE Program, 2017

Canopy Cover


In this study, canopy cover was used to define ecological forestry, intensive forestry, and benign neglect land uses. Because data did not exist for all management areas, a combination of canopy cover certified forest percentage was used as a proxy for these designations.

Map layers were retrieved from the U.S. Geological Survey containing the canopy cover information for the United States (U.S. Geological Survey, 2013). This data was imported and contained to the study region and the pixels were resized to be compatible with the existing map layers. Once the layer had been processed it was combined with both the protection data above as well as the wetland forest data. This final map depicted all the

protection designations as well as the wetland forests that fall within areas with greater than 66% canopy cover (Figure 5).

Figure 5: Wetland forests with greater than 66% canopy cover for each protection designation

Wetland forests under each protection (and potential protection) category where the canopy cover is greater than 66%. For context, wetland forests with less than 66% canopy cover are included.



Source: U.S. Geological Survey, 2014; National Gap Analysis Program, 2015; Snider, 2016; LANDFIRE Program, 2017; U.S. Geological Survey, 2013

To determine the amount of wetland forest converted to the ecological forestry, intensive forestry, and benign neglect land uses, certified forest data was obtained from the Forest Stewardship Council (Forest Stewardship Council, 2016), and the average percent of forests that were certified in the study region was found (2%). This study made the prediction that the area of certified forests would increase in the future (to 5% in the BAU scenario, and 10% in the conservation scenario). The average certified percentage was then applied to the area of wetland forest with greater than 66% canopy cover, and which fell outside of protected areas and SMZs for

each conservation scenario, this area was designated as intensive forestry land use. The remaining unprotected area with greater than 66% canopy cover was designated as benign neglect land use and unprotected wetland forests with canopy cover less than 66% was designated as intensive forestry land use. When applying these values to individual states, the percent of certified forest for each state was used to gain accurate state specific land use areas. For states that currently have large certified percentages (Arkansas - 6.66% & Louisiana - 4.53%) the BAU scenario percentage was raised to 7.5% certified forests by 2050 from the original 5%.

Contextual Information

While the data sets and methods described in the preceding section are sufficient to identify acres of wetland forest for assignment to land use/land management categories and, additional information about these areas provides additional context for the monetary values presented below. Specifically, we present estimates of average tree age and total carbon storage for the areas identified as wetland forests in the first step of our method.

Average Tree Age

Tree age is included in this study to provide an understanding of the average age difference between the protection designations. Average tree age provides some insight into the productivity of the forests, as well as the effectiveness of protection. Country wide data of tree ages was collected and contained to the study region (Pan et. al, 2011). The data was then combined with the wetland forest layers above to produce the age of wetland forests within the study region under each protection designation.

Table 3: Average Age (in Years) of Trees in Current Wetland Forest areas, by Scenario and Land Use Management Group

Scenario:	In Protected Areas	Outside Protected Areas		
		In SMZs	Outside SMZs, Still in Wetland Forests	Converted to Urban Uses (a)
Baseline	45.6	32.9	29.9	0.0
BAU	44.4	32.4	30.2	37.9
Conservation	43.5	30.4	30.6	36.5


Carbon Storage

Carbon sequestration and storage is an important component of the “climate regulation” ecosystem service for which we develop dollar-valued estimates, below. As a side note to that calculation, we have also estimated total carbon storage in all stocks⁴ for the land use/land management categories in each scenario (Table 4). Not that we have not estimated how those carbon stocks may change with shifts of acres among the land use/land management categories. Carbon stores on acres allocated to protected areas, for example, may be assumed to be secure. Stores on acres likely to be developed or managed more intensively for timber would be at risk of losing carbon stocks in the future scenarios.

⁴ These stocks are in live trees (above and below ground), down and standing dead trees, litter, soil organic carbon, and forest understory carbon.

To obtain these estimates, we clipped USDA Forest Service data (Wilson et. al., 2013) to the study region, resampled to 30-meter resolution, and combined the resulting layer with our existing layers defining land use/land management categories (i.e., all wetland forests areas, protected areas by scenario, SMZs by scenario, and wetland forest areas converted to urban uses in the BAU and Conservation scenarios). The estimates do not represent the carbon that would be stored in those categories in the future scenario, but rather the carbon stored in those areas today that may or may not survive the transition to the future land use/land management category. For example, wetland forests converted to urban uses are likely to lose carbon from some stocks in the short run and a diminished capacity to sequester carbon in the long run. Conversely, carbon stocks on areas allocated to protected areas are likely to see their stored carbon remain intact and, perhaps, increase with time. Figure 6 displays carbon stocks per hectare for a small area in eastern North Carolina, and Table 4 presents these estimates in tabular form.

Figure 6: Carbon storage in wetland forests (Mg/ha)



Source: U.S. Geological Survey, 2014; National Gap Analysis Program, 2015; LANDFIRE Program, 2017; Wilson et. al., 2013

Table 4: Carbon Storage (megagrams) in Current Wetland Forests, by Scenario and Land Use/Land Management Group.

Scenario:	In Protected Areas (Mg)	Outside Protected Areas		
		In SMZs (Mg)	Outside SMZs, Still in Wetland Forests (Mg)	Converted to Urban Uses (a) (Mg)
Baseline	162,404,497	70,453,967	1,410,234,595	0
BAU	479,390,228	56,288,738	1,106,753,750	660,343
Conservation	796,375,959	151,866,233	692,261,863	2,589,004

Notes:

a. This column shows estimates of pre-conversion carbon storage in these areas. Carbon storage in areas converted to urban open space and other urban uses will likely be much smaller after conversion.

Methods for Ecosystem Service Valuation

As noted, we use the benefits transfer method, or “BTM,” in which per-unit-area ecosystem service productivity estimates (expressed in dollars per acre per year) from a selection of existing studies of appropriate “source areas” are applied to area estimates in the “policy area.” In the current case, the policy area consists of land that, at least in the baseline case, is wetland forests that is then allocated to various land use/land management categories identified, mapped, or modeled as described above.

Ideally, there would be studies of source areas that provide per-acre estimates of ecosystem service productivity for wetland forests (as distinct from other forests or other wetlands) and, within the wetland forest land use, for different management types (protected, SMZs, in ecological or intensive forestry, etc.). Moreover, one would prefer to have a robust set of these estimates for the full range of ecosystem services of interest in the policy area (see Table 5, above). However, and as noted briefly in the [Executive Summary](#), such a rich set of per-acre ecosystem service values is not available in the literature. There are estimates specific to forests and to wetlands, but few specific to wetland forests.

Moore, et al. (2011), and Schmidt, Moore, and Alber (2014), and Moore (2014), do find that for non-timber ecosystem services, the productivity of wetland forests can be greater than for upland forests. Moreover, it stands to reason that conservation areas, SMZs, and ecological forestry areas that, almost by definition, are managed for a range of ecosystem service values beyond timber, would have higher ecosystem service productivity than areas in intensive, timber-focused forest management. The exception, of course, is the ecosystem service value of raw materials, especially timber, for which intensive forest management would be expected to be more productive.

Table 5. Description of Ecosystem Services Included in the Study

Provisioning Services^a
Food Production: The harvest of agricultural produce, including crops, livestock, and livestock by-products; the food value of hunting, fishing, etc.
Raw Materials: Fuel, fiber, fertilizer, minerals, and energy.
Water Supply: Filtering, retention, storage, and delivery of fresh water—both quality and quantity—for drinking, watering livestock, irrigation, industrial processes, hydroelectric generation, and other uses.
Regulating Services^a
Air Quality: Removing impurities from the air to provide healthy, breathable air for people.
Climate Regulation: Storing atmospheric carbon in biomass and soil as an aid to the mitigation of climate change, and/or keeping regional/local climate (temperature, humidity, rainfall, etc.) within comfortable ranges.
Erosion Control: Retaining arable land, stabilizing slopes, shorelines, riverbanks, etc.
Pollination: Contribution of insects, birds, bats, and other organisms to pollen transport resulting in the production of fruit and seeds. May also include seed and fruit dispersal.
Protection from Extreme Events: Preventing and mitigating impacts on human life, health, and property by attenuating the force of winds, extreme weather events, floods, etc.
Soil Formation: Creation of soil, inducing changes in depth, structure, and fertility, including through nutrient cycling.
Waste Treatment: Improving soil and water quality through the breakdown and/or immobilization of pollution.
Water Flows: Regulation by land cover of the timing of runoff and river discharge, resulting in less severe drought, flooding, and other consequences of too much or too little water available at the wrong time or place.
Cultural Services^a
Aesthetic Value: The role that beautiful, healthy natural areas play in attracting people to live, work, and recreate in a region.
Recreation: The availability of a variety of safe and pleasant landscapes—such as clean water and healthy shorelines—that encourage ecotourism, outdoor sports, fishing, wildlife watching, hunting, etc.

Notes:

a. Descriptions follow Balmford (2010, 2013), Costanza et al. (1997), Reid et al. (2005), and Van der Ploeg, et al. (2010).

These insights suggest the assumptions we employ in this study to bridge the gap between available data and the objective of estimating the ecosystem service value of *wetland* forests in alternative land use/land management categories in our scenarios. Specifically, we make the following assumptions regarding the relative ecosystem service productivity of the various land use/land management classes in our scenarios:

- Protected areas, areas in benign neglect, and SMZs in the conservation scenario (none of which experience active timber management) produce more value per acre per year of most ecosystem services than areas open to timber harvest. The exception is raw material, for which ecosystem service productivity is zero.
- SMZs in the baseline and BAU scenarios, in which timber harvest is possible, will produce less value for most ecosystem services each year than protected (and other no-harvest) areas, but more raw

material/timber value than those areas. Raw material/timber value will be less in SMZs in these scenarios than for areas outside SMZs and available for timber harvest.

- Areas allocated to ecological forestry produce more of most ecosystem service value per acre than areas allocated to intensive forestry or pine plantations. Intensive forestry areas and pine plantations produce more raw material value than areas in ecological forestry.
- No assumptions are necessary for areas converted to urban open space and other urban uses: each of those land use/land management categories has its own set of per-acre ecosystem service productivity estimates.

Table 6, below, embodies these assumptions and states more explicitly, which part of the range of available ecosystem service productivity estimates we have applied in making our estimates of aggregate value per year. We draw the range of candidate ecosystem-service-productivity values from the TEEB database (Van der Ploeg, et al., 2010), from additional studies identified in Phillips and McGee (2014, 2016), and from further literature review undertaken to update those earlier sets of productivity estimates.

Table 6. Rubric for Applying Ecosystem Service Productivity Values, by Land Use/Land Management Category, Scenario, and Ecosystem Service Group.

“Average” means “Average of the range of available values for combinations of ecosystem service and land use.

“Minimum” means the minimum of the same range, and “Maximum” means the maximum of the range. Candidate values for all land use/land management categories except “converted to non-forest use” were drawn from source studies of forests and wetlands. Values for “Converted to non-forest use” are drawn from source studies of urban open space and of other urban land. (Urban open space and Other urban land have separate estimates.)

Land Use / Land Management Category	Ecosystem Service Productivity (value per acres per)					
	Baseline		Business as Usual		Conservation	
	Raw Materials	Other Ecosystem Services	Raw Materials	Other Ecosystem Services	Raw Materials	Other Ecosystem Services
Protected Areas	0	Average	0	Average	0	Average
Added to Protected Areas	0	Average	0	Average	0	Average
Converted to non-forest use	0	Average	0	Average	0	Average
SMZs outside protected areas	Minimum	Average	Minimum	Average	0	Average
Ecological Forestry (outside SMZ)	Average	Average	Average	Average	Average	Average

Intensive Forestry (outside SMZ)	<i>Maximum</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Minimum</i>
Plantation Forestry (outside SMZ)	<i>Maximum</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Minimum</i>	<i>Maximum</i>	<i>Minimum</i>
Benign Neglect	<i>0</i>	<i>Average</i>	<i>0</i>	<i>Average</i>	<i>0</i>	<i>Average</i>

From the full list of available estimates, and keeping with standard BTM practice, we selected individual per-unit-area estimates that came from study areas that are substantially like our policy area. So for example, we eschewed estimates from studies focused on resources in lower-income countries as well as those for exceptionally high income areas in the U.S. We also avoided the oldest studies, some of which were published in the 1970s, and those with estimates that were grossly out of line (and larger than the norm) for similar pairings of land use and ecosystem service. Remaining are 176 estimates of per-acre-per-year ecosystem service value covering all combinations of wetland forest and ecosystem service and, for urban areas, 6 of the 13 ecosystem services. (Where no productivity estimate is available, such as for soil formation in urban areas, no aggregate value estimate is possible, leading to some underestimation of total ecosystem service value for wetland forests that, according to our scenarios, are converted to urban uses.) The full list of the candidate values is provided in [Appendix 2](#).

With the per-acre productivity proxies in hand, aggregate ecosystem service value for each scenario are calculated according to the formula:

$$ESV = \sum_{i,j} [(Acres_j) \times (\$/acre/year)_{i,j}]$$

Where:

$Acres_j$ the number of acres in land use/land management category (j) in the scenario

$(\$/acre/year)_{i,j}$ the dollar value of each ecosystem service (i) provided from each land use (j) each year. These values are drawn from the TEEB database and other sources listed in Appendix 2 and filtered through the rubric described by Table 6, above.

Differences in this aggregate ecosystem service between the baseline and alternative future scenarios can be interpreted as the benefits (since the changes are positive) of changes in land use and land management embodied in those scenarios.

Ecosystem Service Estimates

In our baseline case 35.1 million acres of wetland forests generate \$514.7 billion dollars in ecosystem service value per year (2016\$). (See Tables 7 and 8, below.) Most of these acres and most of this value is in areas currently unprotected from development and open to forest management. In the Business-as-Usual Scenario, in which we assume that half of SECAS-identified high priority conservation lands are placed into protective ownership/management, there is a shift of \$120.7 billion per year in value into the protected areas category. In the Conservation scenario, there is a further shift of acreage, and therefore value, into the protected area category. Thus, by 2050, there could be as much as \$293.5 billion provided by protected wetland forest areas.

Another important driver of differences among the scenarios is the SMZs outside of protected areas. This increase is due to the fact that the SMZs are wider in the conservation scenario. The the increase in value occurs despite the assumption of no commercial harvest within SMZs in that scenario.

Table 7. Estimated Ecosystem Service Value for Wetland Forest in the U.S. South, by Land Use/Land Management Category and Scenario

Land use / Land Management Category	Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:	35,068,824	503,850.0	35,068,824	526,873.0	35,068,824	549,121.6
Protected Areas (GAP 1&2)	2,812,908	51,075.5	9,289,324	169,201.9	15,765,740	287,328.3
Protected in Baseline	2,812,908	51,075.5	2,812,908	51,075.5	2,812,908	51,075.5
Added in Scenario	0	0.0	6,476,416	118,126.4	12,952,832	236,252.8
Converted to Urban Uses	0	0.0	407,334	1,425.7	362,937	1,270.3
Urban Open Space	0	0.0	246,475	828.1	219,611	737.8
Urban Other	0	0.0	160,859	597.6	143,326	532.5
SMZs outside protected areas	1,740,732	31,617.5	1,415,374	25,708.6	3,704,284	67,260.7
SMZs w/in protected areas (a)	70,874	(a)	377,953	(a)	2,216,129	(a)
Available for timber harvest	30,515,184	421,156.9	23,956,791	330,536.8	15,235,862	193,262.3

	Ecological Forestry	453,776	8,276.6	890,188	16,236.6	1,032,274	18,828.1
	Intensive Forestry	7,826,391	9,146.8	5,845,378	6,831.6	4,667,465	5,454.9
	Plantation Forestry	0	0.0	307,651	359.6	245,656	287.1
	Benign Neglect	22,235,017	403,733.5	16,913,573	307,109.1	9,290,466	168,692.1
Notes:							
	a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.						

With regard to the results in Table 7, note that differences between the Baseline and BAU scenarios, and between the Baseline and Conservation scenarios, in the acreage and ecosystem service value for the “converted to urban uses” and “plantation forestry” land use/land management categories does not suggest that conversion of land to those uses increases ecosystem service value. Rather, the increase is an artifact of the baseline situation in which all acres are wetland forests and none of that acreage has yet been converted. Indeed the conversion of wetland forest acres to urban uses does represent a net loss of ecosystem service value relative to the baseline. Acres gained, and value generated on those newly urbanized acres, represent acres lost and (greater) value no longer generated by wetland forests.

In terms of ecosystem services delivered in each scenario, approximately 20% of total ecosystem service value is due to aesthetic value. Protection from extreme events, water flow regulation, and food production supply about 15% each, and water supply and waste treatment contribute up 11.7 and 11.2 percent, respectively, of the total. The balance of ecosystem service value is delivered as air purification, climate regulation, erosion control, pollination, raw materials (timber), recreation, and soil formation. (See Table 8.)

Table 8. Estimated Ecosystem Service Value for Wetland Forest in the U.S. South, by Ecosystem Service and Scenario

	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	503,850.0	526,873.0	549,121.6
Aesthetic Value	101,893.1	106,982.9	111,729.7
Air Quality	4,262.2	4,332.5	4,407.2
Climate Regulation	12,438.8	13,338.4	13,888.2
Erosion Control	1,947.1	2,031.2	2,118.0
Protection from Extreme Events	77,151.4	80,623.6	84,068.8
Food	70,682.5	73,955.5	77,278.3
Pollination	2,079.1	2,172.8	2,268.5
Raw Materials	1,370.8	1,652.9	1,976.5

Recreation	24,135.4	25,689.9	26,778.2
Soil Formation	18,794.9	18,576.5	18,600.3
Waste Treatment	56,215.5	58,810.1	61,446.2
Water Supply	59,135.1	61,981.6	64,738.4
Water Flow Regulation	73,744.0	76,725.2	79,823.4

Conclusions

The study methods and ecosystem service value estimates presented here represent a first approximation of the value currently supplied by wetland forests in the U.S. South. Our model also shows that significant gains in economic value—which at its root reflects better conditions in natural systems—could be obtained through strategies to protect wetland forests already identified as high priorities for conservation and to manage key components (e.g., SMZs) of the majority of wetland forests that will remain part of the region’s working landscape in ways that emphasize and enhance their important ecosystem service contributions.

An important lesson of this effort, however, is that there is much that is not known about the value of wetland forests, particularly in contrast to other forest types and, within the wetland forest type, among different management regimes. Our model has relied on what we regard as reasonable assumptions to develop estimates that are qualitatively sensible, but we would not be surprised to learn from future research that the ecosystem service value of particular areas or even the region as a whole is different than what we have estimated.

The most fruitful path, we imagine, for next research steps would involve downscaling our modeling approach to utilize spatial information on sub-classes of wetland forests, such as mangroves, or on smaller areas for which per-acre ecosystem service productivity would be influenced by proximity to population centers, infrastructure, and/or other natural resources for which wetland forests provide more or less of different ecosystem services than the broad ranges and averages employed in our method would suggest.

Meanwhile, it is our hope that this study will provide, at minimum, a means of advancing conversations about both how to ensure the future of wetland forest and the human values that depend on them and how to advance essential knowledge about the magnitude of those values for planning purposes.

Works Cited

- Aburto-Oropeza, O., Ezcurra, E., Danemann, G., Valdez, V., Murray, J., & Sala, E. (2008). Mangroves in the Gulf of California increase fishery yields. *Proceedings of the National Academy of Sciences of the United States of America*, 105(30), 10456–10459. <https://doi.org/10.1073/pnas.0804601105>
- Amacher, G. S., & Brazee, R. J. (1989). Application of wetland valuation techniques: Examples from Great Lakes Coastal wetlands. *University of Michigan, School of Natural Resources*.
- Anderson, G. D., & Edwards, S. F. (1986). Protecting Rhode Island coastal salt ponds - an economic-assessment of downzoning. *Coastal Zone Management Journal*, 14, 67–91.

Balmford, A., Fisher, B., Green, R. E., Naidoo, R., Strassburg, B., Kerry Turner, R., & Rodrigues, A. S. L. (2010). Bringing Ecosystem Services into the Real World: An Operational Framework for Assessing the Economic Consequences of Losing Wild Nature. *Environmental and Resource Economics*, 48(2), 161–175. <https://doi.org/10.1007/s10640-010-9413-2>

Balmford, A., Rodrigues, A., Walpole, M., Brink, P., Kettunen, M., de Groot, R., & Cambridge, U. (2013). The Economics of Biodiversity and Ecosystems: Scoping the Science. (Vol. 8 SRC-GoogleScholar). Retrieved from http://ec.europa.eu/environment/nature/biodiversity/economics/teeb_en.htm

Batie, S. S., & Wilson, J. R. (1978). Economic Values Attributable to Virginia's Coastal Wetlands as Inputs in Oyster Production. *Southern Journal of Agricultural Economics*. Retrieved from <http://ageconsearch.umn.edu/bitstream/30270/1/10010111.pdf>

Bell, F. W. (1989). *Application of wetland valuation theory to commercial and recreational fisheries in Florida* (p. 148). Florida Sea Grant Program.

Belyea, C. M., & Terando, A. J. (2014). Designing Sustainable Landscapes (SLEUTH). Biodiversity and Spatial Information Center: NC State University. Retrieved from <http://www.basic.ncsu.edu/dsl/urb.html>

Bergstrom, J. C., Stoll, J. R., Titre, J. P., & Wright, V. L. (1990). Economic value of wetlands-based recreation. *Ecological Economics*, 2(2), 129–147. [https://doi.org/10.1016/0921-8009\(90\)90004-E](https://doi.org/10.1016/0921-8009(90)90004-E)

Breaux, A., Farber, S., & Day, J. (1995). Using Natural Coastal Wetlands Systems for Wastewater Treatment: An Economic Benefit Analysis. *Journal of Environmental Management*, 44(3), 285–291. <https://doi.org/10.1006/jema.1995.0046>

Brenner Guillermo, J. (2007, May 4). Valuation of ecosystem services in the catalan coastal zone [info:eu-repo/semantics/doctoralThesis]. Retrieved May 18, 2014, from <http://www.tdx.cat/handle/10803/6398>

Byström, O. (2000). The Replacement Value of Wetlands in Sweden. *Environmental and Resource Economics*, 16(4), 347–362. <https://doi.org/10.1023/A:1008316619355>

Costanza, R., d'Arge, R., Farber, S., Grasso, M., deGroot, R., Hannon, B., & van den Belt, M. (1997). The Value of the World's Ecosystem Services and Natural Capital. *Nature*, 387, 253–260.

Costanza, R., d'Arge, R., Farber, S., Grasso, M., deGroot, R., Hannon, B., & van den Belt, M. (1997). The Value of the World's Ecosystem Services and Natural Capital. *Nature*, 387, 253–260.

Costanza, Robert, & Farley, J. (2007). Ecological economics of coastal disasters: Introduction to the special issue. *Ecological Economics*, 63(2–3), 249–253. <https://doi.org/10.1016/j.ecolecon.2007.03.002>

Costanza, Robert, Farber, S. C., & Maxwell, J. (1989). Valuation and management of wetland ecosystems. *Ecological Economics*, 1(4), 335–361. [https://doi.org/10.1016/0921-8009\(89\)90014-1](https://doi.org/10.1016/0921-8009(89)90014-1)

Costanza, Robert, Wilson, M., Troy, A., Voinov, A., Liu, S., & D'Agostino, J. (2006). The value of New Jersey's ecosystem services and natural capital. *Gund Institute for Ecological Economics, University of Vermont and New Jersey Department of Environmental Protection, Trenton, New Jersey, 13*. Retrieved from <http://www.academia.edu/download/30561335/njvaluationpart2.pdf>

Creel, M., & Loomis, J. (1992). Recreation value of water to wetlands in the San Joaquin Valley: Linked multinomial logit and count data trip frequency models. *Water Resources Research*, 28(10), 2597–2606. <https://doi.org/10.1029/92WR01514>

Cruz, A. de la, & Benedicto, J. (2009). *Assessing Socio-economic Benefits of Natura 2000 – a Case Study on the ecosystem service provided by SPA PICO DA VARA / RIBEIRA DO GUILHERME*. (Output of the project Financing Natura 2000: Cost estimate and benefits of Natura 2000 (Contract No.: 070307/2007/484403/MAR/B2).). 43. Retrieved from http://ec.europa.eu/environment/nature/natura2000/financing/docs/azores_case_study.pdf

Dahl, T. E. (2011). *Status and Trends of Wetlands in the Conterminous United States 2004 to 2009* (Report to Congress). U.S. Department of the Interior: U.S. Fish and Wildlife Service. Retrieved from <https://www.fws.gov/wetlands/documents/Status-and-Trends-of-Wetlands-in-the-Conterminous-United-States-2004-to-2009.pdf>

Everard, M., Great Britain, & Environment Agency. (2009). *Ecosystem services case studies*. Bristol: Environment Agency.

Farley, J. (2012). Ecosystem services: The economics debate. *Ecosystem Services*, 1(1), 40–49. <https://doi.org/10.1016/j.ecoser.2012.07.002>

Flores, L., Harrison-Cox, J., Wilson, S., & Batker, D. (2013). *Nearshore Valuation-Primary Values. Nature's Value in Clallam County: The Economic Benefits of Feeder Bluffs and 12 Other Ecosystems*. Tacoma, WA: Earth Economics.

Folke, C., & Kaberger. (1991). The societal value of wetland life-support. In C. Folke & T. Kåberger (Eds.), *Linking the natural environment and the economy*. Dordrecht: Springer Netherlands. Retrieved from <http://link.springer.com/10.1007/978-94-017-6406-3>

Forest Stewardship Council. (2016). Facts & Figures: FSC Certified Acres by State. Retrieved July 17, 2017, from <https://us.fsc.org:443/en-us/what-we-do/facts-figures>

Fry, J., G. Xian, S. Jin, J. Dewitz, C. Homer, L. Yang, C. Barnes, N. Herold, and J. Wickham. (2011). "NLCD_September2011PERS.pdf." *Photogrammetric Engineering & Remote Sensing* 77 (9): 858–64.

Gerrans, P. (1994). An economic valuation of the Jandakot wetlands. *ECU Publications Pre. 2011*. Retrieved from <http://ro.ecu.edu.au/ecuworks/6867>

- Gren, I.-M., & Söderqvist, T. (1994). *Economic valuation of wetlands: a survey*. Beijer International Institute of Ecological Economics, The Royal Swedish Academy of Sciences.
- Gupta, T. R., & Foster, J. H. (1975). Economic criteria for freshwater wetland policy in Massachusetts. *American Journal of Agricultural Economics*. Retrieved from <http://agris.fao.org/agris-search/search.do?recordID=US19750030691>
- Haener, M. K., & Adamowicz, W. L. (2000). Regional forest resource accounting: a northern Alberta case study. *Canadian Journal of Forest Research*, 30(2), 264–273. <https://doi.org/10.1139/x99-213>
- Hall, D. (2010, December). “Applied ecosystem services in working forests: A direct market valuation” by Darin Stuart Hale. University of Tennessee, Knoxville. Retrieved from http://trace.tennessee.edu/utk_gradthes/804/
- Hamilton, L. S., & Snedaker, S. C. (1984). *Handbook for mangrove area management*. Honolulu: East-West Environment and Policy Institute. Retrieved from <http://scholarspace.manoa.hawaii.edu/handle/10125/26547>
- Hughes, Z. (2006). *Ecological and economics assessment of potential eelgrass expansion Hughesseat Sucia Island* (Term Paper). University of Washington.
- Jaworski, E., & Raphael, C. N. (1978). Fish, Wildlife, And Recreational Values of Michigan 's Coastal Wetlands. *Michigan Department of Natural Resources February 1978, 209 P. 25 Fig, 62 Tab, 205 Ref, 1 Append. Wetlands Value Study, Phase 1*. Retrieved from <http://search.proquest.com/docview/19192481?pq-origsite=summon&>
- Jenkins, W. A., Murray, B. C., Kramer, R. A., & Faulkner, S. P. (2010). Valuing ecosystem services from wetlands restoration in the Mississippi Alluvial Valley. *Ecological Economics*, 69(5), 1051–1061. <https://doi.org/10.1016/j.ecolecon.2009.11.022>
- Johnston, R. J., Opaluch, J. J., Grigalunas, T. A., & Mazzotta, M. J. (2001). Estimating Amenity Benefits of Coastal Farmland. *Growth and Change*, 32(3), 305–325. <https://doi.org/10.1111/0017-4815.00161>
- Kauffman, G., Homsey, A., McVey, E., Mack, S., & Chatterson, S. (2011). *Socioeconomic Value of the Chesapeake Bay Watershed in Delaware* (p. 45). Newark, Delaware: University of Delaware's Institute for Public Administration. Retrieved from <http://www.wra.udel.edu/wp-content/publications/DelChesapeakeWatershed.pdf>
- Kosz, M., Brezina, B., & Madreiter, T. (1992). *Kosten-Nutzen-Analyse ausgewählter Varianten eines Nationalparks Donauauen*. Institut für Finanzwissenschaft und Infrastrukturpolitik der Technischen Universität Wien.
- Kreutzwiser, R. (1981). The Economic Significance of the Long Point Marsh, Lake Erie, as a Recreational Resource. *Journal of Great Lakes Research*, 7(2), 105–110. [https://doi.org/10.1016/S0380-1330\(81\)72034-3](https://doi.org/10.1016/S0380-1330(81)72034-3)

LANDFIRE Program. (2017). *LANDFIRE Program: Data Products - Vegetation - Existing Vegetation Type* (Data). USDA Forest Service and US Department of the Interior. Retrieved from <https://www.landfire.gov/NationalProductDescriptions21.php>

Lant, C., & Roberts, R. (1990). Greenbelts in the Cornbelt: Riparian Wetlands, Intrinsic Values and Market Failure. *Environment and Planning A*, 1375–1388.

Leschine, T. M., Wellman, K. F., & Green, T. H. (1997). *The Economic Value of Wetlands: Wetlands' Role in Flood Protection in Western Washington* (Ecology Publication No. 97-100). Washington State Department of Ecology. Retrieved from <https://fortress.wa.gov/ecy/publications/publications/97100.pdf>

Loomis, J., & Ekstrand, E. (1998). Alternative approaches for incorporating respondent uncertainty when estimating willingness to pay: the case of the Mexican spotted owl. *Ecological Economics*, 27(1), 29–41.

Lui, Z. (2006). *Water Quality Simulation and Economic Valuation of Riparian Land-use Changes*. University of Cincinnati.

Luisetti, T., Turner, R. K., Bateman, I. J., Morse-Jones, S., Adams, C., & Fonseca, L. (2011). Coastal and marine ecosystem services valuation for policy and management: Managed realignment case studies in England. *Ocean & Coastal Management*, 54(3), 212–224. <https://doi.org/10.1016/j.ocecoaman.2010.11.003>

Mates, W. (2007). *Valuing New Jersey's Natural Capital: An Assessment of the Economic Value of the State's Natural Resources* (p. 54). New Jersey Department of Environmental Protection, Division of Science, Research, and Technology. Retrieved from <http://www.state.nj.us/dep/dsr/naturalcap/nat-cap-1.pdf>

Mazzotta, M. J. (1996). Measuring public values and priorities for natural resources: An application to the Peconic Estuary system. *Dissertations and Master's Theses (Campus Access)*, 1–272.

McPherson, G. E. (1992). Accounting for benefits and costs of urban greenspace. *Landscape and Urban Planning*, 22(1), 41–51. [https://doi.org/10.1016/0169-2046\(92\)90006-L](https://doi.org/10.1016/0169-2046(92)90006-L)

McPherson, G., Scott, K., & Simpson, J. (1998). Estimating cost effectiveness of residential yard trees for improving air quality in Sacramento, California, using existing models. *Atmospheric Environment*, 32(1), 75–84. [https://doi.org/10.1016/S1352-2310\(97\)00180-5](https://doi.org/10.1016/S1352-2310(97)00180-5)

Moore, R., Williams, T., Rodriguez, E., & Hepinstall-Cymmerman, J. (2011). *Quantifying the value of non-timber ecosystem services from Georgia's private forests* (p. 44). Retrieved from <http://www.warnell.uga.edu/news/wp-content/uploads/2011/02/Final-Report-1-24-11.pdf>

Moore, R., Williams, T., Rodriguez, E., & Hepinstall-Cymmerman, J. (2011). *Quantifying the value of non-timber ecosystem services from Georgia's private forests* (p. 44). Retrieved from <http://www.warnell.uga.edu/news/wp-content/uploads/2011/02/Final-Report-1-24-11.pdf>

- Morton, R. M. (1990). Community structure, density and standing crop of fishes in a subtropical Australian mangrove area. *Marine Biology*, 105(3), 385–394. <https://doi.org/10.1007/BF01316309>
- National Gap Analysis Program. (2015). National Gap Analysis Program: USGS Core Sciences Analytics and Synthesis. U.S. Geological Survey. Retrieved from <http://gapanalysis.usgs.gov/>
- Nowak, D. J., Crane, D. E., Dwyer, J. F., & others. (2002). Compensatory value of urban trees in the United States. *Journal of Arboriculture*, 28(4), 194–199.
- NRDC. 2015. “In the U.S. Southeast, Natural Forests Are Being Felled to Send Fuel Overseas.” Natural Resources Defense Council.
- Opaluch, J. J., Grigalunas, T. A., Diamantedes, J., Mazzotta, M., & Johnston, R. (1999). *Recreational and resource economic values for the Peconic Estuary, prepared for the Peconic Estuary Program*. Peconic Estuary Program.
- Pan, Y., Chen, J. M., Birdsey, R., McCullough, K., He, L., & Deng, F. (2011). Age structure and disturbance legacy of North American forests. *Biogeosciences*, 8, 715–732. <https://doi.org/10.5194/bg-8-715-2011>
- Prince, R., & Ahmed, E. (1989). Estimating individual recreation benefits under congestion and uncertainty. *Journal of Leisure Research*, 21, 61–76.
- Qiu, Z., Prato, T., & Boehrn, G. (2006). Economic Valuation of Riparian Buffer and Open Space in a Suburban Watershed1. *JAWRA Journal of the American Water Resources Association*, 42(6), 1583–1596. <https://doi.org/10.1111/j.1752-1688.2006.tb06022.x>
- Rausser, G. C., & Small, A. A. (2000). *Valuing Research Leads: Bioprospecting and the Conservation of Genetic Resources* (Berkeley Olin Program in Law & Economics, Working Paper Series). Berkeley Olin Program in Law & Economics. Retrieved from <http://econpapers.repec.org/paper/cdloplwec/qt4t56m5b8.htm>
- Reid, W. V., Mooney, H. A., Cooper, A., Capistrano, D., Carpenter, S. R., Chopra, K., ... Zurek, M. B. (2005). Millennium Ecosystem Assessment, Ecosystems and Human Well-being: Synthesis. Washington, DC: Island Press.
- Schmidt, J. P., Moore, R., & Alber, M. (2014). Integrating ecosystem services and local government finances into land use planning: A case study from coastal Georgia. *Landscape and Urban Planning*, 122, 56–67. <https://doi.org/10.1016/j.landurbplan.2013.11.008>
- Shafer, E. L., Carline, R., Guldin, R. W., & Cordell, H. K. (1993). Economic amenity values of wildlife: Six case studies in Pennsylvania. *Environmental Management*, 17(5), 669–682. <https://doi.org/10.1007/BF02393728>
- Snider, M. (2016). SECAS Blueprint v1.0 | Southeast Region CPA. Retrieved March 28, 2017, from <https://seregon.databasin.org/datasets/c237c946ba0c49b0957cffe1bfb7ad7>

Streiner, C. F., & Loomis, J. B. (1995). Estimating the Benefits of Urban Stream Restoration Using the Hedonic Price Method.

T.r, G., & J.h, F. (1975). Economic criteria for freshwater wetland policy in Massachusetts. *American Journal of Agricultural Economics*. Retrieved from <http://agris.fao.org/agris-search/search.do?recordID=US19750030691>

The Trust for Public Land. (2010). *The economic benefits and fiscal impact of parks and open space in Nassau and Suffolk Counties, New York* (p. 48). The Trust for Public Land. Retrieved from <http://cloud.tpl.org/pubs/ccpe--nassau-county-park-benefits.pdf>

Thibodeau, F. R., & Ostro, B. D. (1981). An economic analysis of wetland protection. *Journal of Environmental Management*, 19, 72–79.

U.S. Geological Survey. (2013). Tree Canopy Data For Tree Canopy - conterminous Unites states....100 meter resolution. Retrieved from https://nationalmap.gov/small_scale/atlasftp.html?openChapters=chpbio#chpbio

U.S. Geological Survey. (2014). *NLCD 2011 Land Cover (2011 Edition, amended 2014) - National Geospatial Data Asset (NGDA) Land Use Land Cover*. Sioux Falls, SD: U.S. Geological Survey. Retrieved from <http://www.mrlc.gov/nlcd2006.php>

U.S. Geological Survey. (2017). U.S. Geological Survey - National Hydrography Dataset. Retrieved July 17, 2017, from https://nhd.usgs.gov/NHD_High_Resolution.html

UK Environment Agency. (1999). *River Ancholme flood storage area progression*. (No. E3475/01/001). Prepared by Posford Duvivier Environment.

Van der Ploeg, S., Wang, Y., Gebre Weldmichael, T., & De Groot, R. S. (2010). *The TEEB Valuation Database – a searchable database of 1310 estimates of monetary values of ecosystem services*. (Excel database and accompanying documentation). Wageningen, The Netherlands: Foundation for Sustainable Development. Retrieved from <http://www.es-partnership.org/esp/80763/5/0/50>

Weber, T. (2007). *Ecosystem services in Cecil County's green infrastructure: Technical Report for the Cecil County Green Infrastructure Plan* (White Paper) (p. 32). Annapolis, MD: The Conservation Fund. Retrieved from http://www.ccgov.org/uploads/PlanningAndZoning/General/CecilCoMD_TechReport%20-%20Ecosystem%20services.pdf

Whitehead, J. C. (1990). Measuring willingness-to-pay for wetlands preservation with the contingent valuation method. *Wetlands*, 10(2), 187–201. <https://doi.org/10.1007/BF03160832>

Whitehead, J. C., Groothuis, P. A., Southwick, R., & Foster-Turley, P. (2009). Measuring the economic benefits of Saginaw Bay coastal marsh with revealed and stated preference methods. *Journal of Great Lakes Research*, 35(3), 430–437. <https://doi.org/10.1016/j.jglr.2009.03.005>

Wilson, B. T., Woddall, C. W., & Griffith, D. M. (2013). *Forest carbon stocks of the contiguous United States (2000-2009)*. Newtown Square, PA: USDA Forest Service, Northern Research Station. Retrieved from <http://www.fs.usda.gov/rds/archive/Product/RDS-2013-0004>

Zhou, X., Al-Kaisi, M., & Helmers, M. J. (2009). Cost effectiveness of conservation practices in controlling water erosion in Iowa. *Soil and Tillage Research*, 106(1), 71–78. <https://doi.org/10.1016/j.still.2009.09.015>

APPENDIX 1: State-by-State Estimates

(Tables begin next page)

For each of 14 states, Table xx-1 (where “xx” is the state abbreviation) shows acreage and ecosystem service value for each land use-land management category and for each scenario (baseline, business-as-usual, and conservation). Table xx-2 shows ecosystem service value, by ecosystem service, for each scenario. All dollar values reflect 2016 price levels, and all dollar-values are values generated per year.

Alabama

Table AL-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Alabama

Land use / Land Management Category			Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
			Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:			2,196,499	34,370	2,196,499	35,522	2,196,499	36,660
	Protected Areas (GAP 1&2)		90,608	1,645	589,967	10,753	1,089,327	19,861
	Protected in Baseline		90,608	1,645	90,608	1,645	90,608	1,645
	Added in Scenario		0	0	499,360	9,108	998,719	18,216
	Converted to Urban Uses		0	0	9,233	32	8,108	28
	Urban Open Space		0	0	5,587	19	4,906	16
	Urban Other		0	0	3,646	14	3,202	12
	SMZs outside protected areas		138,055	2,507	111,085	2,018	317,625	5,767
	SMZs w/in protected areas (a)		1,982	(a)	28,281	(a)	211,097	(a)
	Available for timber harvest		1,967,835	30,218	1,486,213	22,719	781,439	11,003
	Ecological Forestry		45,514	830	61,738	1,126	59,359	1,083
	Intensive Forestry		324,736	380	238,875	279	178,453	209
	Plantation Forestry		0	0	12,572	15	9,392	11
	Benign Neglect		1,597,586	29,008	1,173,027	21,299	534,234	9,700
Notes:								
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.								

Table AL-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Alabama.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	34,370	35,522	36,660
Aesthetic Value	6,999	7,246	7,486
Air Quality	277	280	284
Climate Regulation	854	891	920
Erosion Control	133	137	142
Protection from Extreme Events	5,277	5,451	5,625
Food	4,855	5,021	5,188
Pollination	143	147	152
Raw Materials	59	89	118
Recreation	1,658	1,725	1,781
Soil Formation	1,177	1,172	1,173
Waste Treatment	3,861	3,992	4,125
Water Supply	4,061	4,201	4,341
Water Flow Regulation	5,017	5,170	5,326

Arkansas

Table AR-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Arkansas

Land use / Land Management Category	Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:	2,491,063	37,430	2,491,063	38,852	2,491,063	40,272
Protected Areas (GAP 1&2)	459,387	8,341	891,769	16,228	1,324,152	24,114
Protected in Baseline	459,387	8,341	459,387	8,341	459,387	8,341
Added in Scenario	0	0	432,383	7,886	864,765	15,773
Converted to Urban Uses	0	0	3,126	11	3,006	11
Urban Open Space	0	0	1,892	6	1,819	6
Urban Other	0	0	1,235	5	1,187	4
SMZs outside protected areas	168,937	3,069	137,079	2,490	248,979	4,521
SMZs w/in protected areas (a)	17,268	(a)	48,798	(a)	196,834	(a)
Available for timber harvest	1,862,739	26,020	1,459,088	20,124	914,925	11,626
Ecological Forestry	93,439	1,704	81,282	1,483	62,111	1,133
Intensive Forestry	459,759	537	356,561	417	279,123	326
Plantation Forestry	0	0	18,766	22	14,691	17
Benign Neglect	1,309,542	23,778	1,002,479	18,203	559,001	10,150
Notes:						
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.						

Table AR-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Arkansas.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	s	38,852	40,272
Aesthetic Value	7,596	7,902	8,206
Air Quality	308	313	318
Climate Regulation	927	967	1,004
Erosion Control	145	150	156
Protection from Extreme Events	5,739	5,958	6,177
Food	5,270	5,480	5,691
Pollination	155	161	167
Raw Materials	86	106	126
Recreation	1,799	1,875	1,947
Soil Formation	1,335	1,333	1,333
Waste Treatment	4,191	4,357	4,525
Water Supply	4,408	4,584	4,760
Water Flow Regulation	5,470	5,665	5,861

Florida

Table FL-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Florida

Land use / Land Management Category			Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
			Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:			5,361,255	80,797	5,361,255	84,664	5,361,255	88,306
	Protected Areas (GAP 1&2)		665,867	12,091	1,776,665	32,351	2,887,464	52,611
		Protected in Baseline	665,867	12,091	665,867	12,091	665,867	12,091
		Added in Scenario	0	0	1,110,798	20,260	2,221,597	40,521
	<i>Converted to Urban Uses</i>		0	0	100,699	352	87,675	307
		Urban Open Space	0	0	60,932	205	53,052	178
		Urban Other	0	0	39,767	148	34,624	129
	<i>SMZs outside protected areas</i>		87,936	1,597	66,953	1,216	196,376	3,566
	<i>SMZs w/in protected areas (a)</i>		7,616	(a)	25,894	(a)	183,717	(a)
	<i>Available for timber harvest</i>		4,607,453	67,109	3,416,938	50,745	2,189,740	31,822
		Ecological Forestry	22,162	404	137,562	2,509	172,164	3,140
		Intensive Forestry	974,316	1,139	632,413	739	444,693	520
		Plantation Forestry	0	0	33,285	39	23,405	27
		Benign Neglect	3,610,974	65,566	2,613,678	47,458	1,549,478	28,135
Notes:								
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.								

Table FL-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Florida.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	80,797	84,664	88,306
Aesthetic Value	16,405	17,272	18,045
Air Quality	664	676	688
Climate Regulation	2,003	2,177	2,263
Erosion Control	313	326	341
Protection from Extreme Events	12,392	12,968	13,532
Food	11,381	11,918	12,463
Pollination	334	350	365
Raw Materials	167	215	275
Recreation	3,886	4,178	4,350
Soil Formation	2,873	2,819	2,826
Waste Treatment	9,050	9,476	9,908
Water Supply	9,520	9,997	10,447
Water Flow Regulation	11,809	12,293	12,802

Georgia

Table GA-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Georgia

Land use / Land Management Category	Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:	4,371,809	67,313	4,371,809	68,818	4,371,809	70,096
Protected Areas (GAP 1&2)	380,048	6,901	895,236	16,297	1,410,423	25,694
Protected in Baseline	380,048	6,901	380,048	6,901	380,048	6,901
Added in Scenario	0	0	515,187	9,397	1,030,374	18,793
Converted to Urban Uses	0	0	104,518	366	93,556	327
Urban Open Space	0	0	63,243	212	56,610	190
Urban Other	0	0	41,275	153	36,946	137
SMZs outside protected areas	124,486	2,261	105,636	1,919	608,374	11,047
SMZs w/in protected areas (a)	2,228	(a)	18,186	(a)	229,345	(a)
Available for timber harvest	3,867,275	58,152	3,266,420	50,236	2,259,457	33,028
Ecological Forestry	12,943	236	136,583	2,491	178,778	3,261
Intensive Forestry	710,443	830	508,026	594	448,090	524
Plantation Forestry	0	0	26,738	31	23,584	28
Benign Neglect	3,143,889	57,085	2,595,073	47,120	1,609,005	29,216
Notes:						
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for						

Protected Areas. Acreage is provided here for reference.

Table GA-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Georgia.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	67,313	68,818	70,096
Aesthetic Value	13,691	14,051	14,317
Air Quality	547	550	554
Climate Regulation	1,671	1,787	1,812
Erosion Control	261	265	270
Protection from Extreme Events	10,331	10,540	10,737
Food	9,498	9,681	9,873
Pollination	279	284	290
Raw Materials	122	144	179
Recreation	3,243	3,418	3,472
Soil Formation	2,343	2,287	2,293
Waste Treatment	7,552	7,697	7,850
Water Supply	7,944	8,127	8,284
Water Flow Regulation	9,832	9,986	10,166

Kentucky

Table KY-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Kentucky

Land use / Land Management Category			Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
			Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:			179,959	1,911	179,959	2,155	179,959	2,399
	Protected Areas (GAP 1&2)		16,211	294	49,201	896	82,191	1,498
		Protected in Baseline	16,211	294	16,211	294	16,211	294
		Added in Scenario	0	0	32,990	602	65,980	1,203
	<i>Converted to Urban Uses</i>		0	0	1,237	4	955	3
		Urban Open Space	0	0	749	3	578	2
		Urban Other	0	0	489	2	377	1
	<i>SMZs outside protected areas</i>		14,968	272	11,724	213	22,604	410
	SMZs w/in protected areas (a)		944	(a)	4,038	(a)	18,988	(a)
	<i>Available for timber harvest</i>		148,780	1,344	117,796	1,042	74,209	487
		Ecological Forestry	1,385	25	2,660	49	2,355	43
		Intensive Forestry	79,897	93	61,363	72	48,126	56
		Plantation Forestry	0	0	3,230	4	2,533	3
		Benign Neglect	67,498	1,226	50,543	918	21,195	385
Notes:								
	a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.							

Table KY-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Kentucky.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	1,911	2,155	2,399
Aesthetic Value	375	428	481
Air Quality	20	20	21
Climate Regulation	46	53	59
Erosion Control	7	8	9
Protection from Extreme Events	289	327	365
Food	260	296	333
Pollination	8	9	10
Raw Materials	14	14	14
Recreation	89	102	115
Soil Formation	96	96	96
Waste Treatment	207	236	265
Water Supply	218	248	279
Water Flow Regulation	283	316	351

Louisiana

Table LA-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Louisiana

Land use / Land Management Category	Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:	5,231,118	79,426	5,231,118	82,700	5,231,118	85,959
Protected Areas (GAP 1&2)	244,379	4,437	1,422,930	25,933	2,601,482	47,430
Protected in Baseline	244,379	4,437	244,379	4,437	244,379	4,437
Added in Scenario	0	0	1,178,551	21,496	2,357,103	42,992
<i>Converted to Urban Uses</i>	<i>0</i>	<i>0</i>	<i>6,580</i>	<i>23</i>	<i>5,738</i>	<i>20</i>
Urban Open Space	0	0	3,982	13	3,472	12
Urban Other	0	0	2,599	10	2,266	8
<i>SMZs outside protected areas</i>	<i>242,374</i>	<i>4,402</i>	<i>185,232</i>	<i>3,364</i>	<i>301,813</i>	<i>5,480</i>
SMZs w/in protected areas (a)	14,782	(a)	71,362	(a)	290,402	(a)
<i>Available for timber harvest</i>	<i>4,744,365</i>	<i>70,587</i>	<i>3,616,375</i>	<i>53,380</i>	<i>2,322,086</i>	<i>33,029</i>
Ecological Forestry	173,394	3,163	216,916	3,956	178,356	3,253
Intensive Forestry	916,673	1,071	687,959	804	511,597	598
Plantation Forestry	0	0	36,208	42	26,926	31
Benign Neglect	3,654,297	66,353	2,675,292	48,577	1,605,206	29,147
Notes:						
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.						

Table LA-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Louisiana.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	79,426	82,700	85,959
Aesthetic Value	16,134	16,833	17,527
Air Quality	650	661	672
Climate Regulation	1,970	2,060	2,144
Erosion Control	307	320	332
Protection from Extreme Events	12,183	12,684	13,185
Food	11,192	11,673	12,156
Pollination	329	343	357
Raw Materials	170	238	299
Recreation	3,822	3,993	4,157
Soil Formation	2,804	2,800	2,801
Waste Treatment	8,900	9,282	9,664
Water Supply	9,362	9,765	10,167
Water Flow Regulation	11,604	12,050	12,499

Mississippi

Table MS-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Mississippi

Land use / Land Management Category		Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
		Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:		3,502,004	51,044	3,502,004	53,169	3,502,004	55,239
Protected Areas (GAP 1&2)		172,543	3,133	897,343	16,353	1,622,142	29,573
	Protected in Baseline	172,543	3,133	172,543	3,133	172,543	3,133
	Added in Scenario	0	0	724,799	13,220	1,449,598	26,440
<i>Converted to Urban Uses</i>		0	0	25,864	91	22,019	77
	Urban Open Space	0	0	15,650	53	13,324	45
	Urban Other	0	0	10,214	38	8,696	32
<i>SMZs outside protected areas</i>		164,550	2,989	134,712	2,447	467,321	8,485
SMZs w/in protected areas (a)		3,956	(a)	32,549	(a)	278,565	(a)
<i>Available for timber harvest</i>		3,164,910	44,923	2,444,085	34,278	1,390,522	17,104
	Ecological Forestry	37,123	677	92,455	1,686	91,067	1,661
	Intensive Forestry	738,564	863	565,226	661	455,857	533
	Plantation Forestry	0	0	29,749	35	23,992	28
	Benign Neglect	2,389,223	43,382	1,756,654	31,897	819,605	14,882
Notes:							
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.							

Table MS-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Mississippi.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	51,044	53,169	55,239
Aesthetic Value	10,335	10,798	11,238
Air Quality	428	435	442
Climate Regulation	1,262	1,336	1,387
Erosion Control	197	205	213
Protection from Extreme Events	7,820	8,141	8,460
Food	7,170	7,474	7,782
Pollination	211	220	228
Raw Materials	129	168	207
Recreation	2,448	2,580	2,681
Soil Formation	1,877	1,863	1,865
Waste Treatment	5,702	5,943	6,188
Water Supply	5,998	6,259	6,515
Water Flow Regulation	7,468	7,747	8,034

Missouri

Table MO-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Missouri

Land use / Land Management Category	Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:	692,427	7,736	692,427	8,362	692,427	8,986
Protected Areas (GAP 1&2)	103,782	1,884	191,341	3,481	278,899	5,078
Protected in Baseline	103,782	1,884	103,782	1,884	103,782	1,884
Added in Scenario	0	0	87,558	1,597	175,117	3,194
<i>Converted to Urban Uses</i>	0	0	1,216	4	1,176	4
Urban Open Space	0	0	736	2	711	2
Urban Other	0	0	480	2	464	2
<i>SMZs outside protected areas</i>	90,576	1,646	81,212	1,476	175,533	3,187
SMZs w/in protected areas (a)	5,316	(a)	14,551	(a)	63,686	(a)
<i>Available for timber harvest</i>	498,069	4,206	418,658	3,401	236,819	716
Ecological Forestry	64	1	8,567	156	2,582	47
Intensive Forestry	284,763	333	234,950	275	200,446	234
Plantation Forestry	0	0	12,366	14	10,550	12
Benign Neglect	213,243	3,872	162,776	2,956	23,241	422
Notes:						
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.						

Table MO-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Missouri.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	7,736	8,362	8,986
Aesthetic Value	1,526	1,663	1,798
Air Quality	77	79	81
Climate Regulation	186	204	220
Erosion Control	30	32	35
Protection from Extreme Events	1,173	1,271	1,369
Food	1,059	1,152	1,246
Pollination	31	34	37
Raw Materials	49	51	50
Recreation	361	395	427
Soil Formation	371	370	370
Waste Treatment	843	917	992
Water Supply	887	965	1,044
Water Flow Regulation	1,143	1,230	1,317

North Carolina

Table NC-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for North Carolina

Land use / Land Management Category			Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
			Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:			2,415,774	26,939	2,415,774	29,526	2,415,774	32,063
	Protected Areas (GAP 1&2)		161,613	2,935	520,080	9,473	878,548	16,011
		Protected in Baseline	161,613	2,935	161,613	2,935	161,613	2,935
		Added in Scenario	0	0	358,467	6,538	716,935	13,077
	Converted to Urban Uses		0	0	33,857	118	30,092	105
		Urban Open Space	0	0	20,487	69	18,209	61
		Urban Other	0	0	13,370	50	11,884	44
	SMZs outside protected areas		165,978	3,015	140,464	2,552	324,789	5,897
	SMZs w/in protected areas (a)		2,166	(a)	25,285	(a)	134,920	(a)
	Available for timber harvest		2,088,182	20,989	1,721,372	17,383	1,182,345	10,049
		Ecological Forestry	9,280	169	45,229	825	50,993	930
		Intensive Forestry	996,395	1,164	775,953	907	638,796	747
		Plantation Forestry	0	0	40,840	48	33,621	39
		Benign Neglect	1,082,507	19,656	859,350	15,604	458,935	8,333
Notes:								
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.								

Table NC-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, North Carolina.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	26,939	29,526	32,063
Aesthetic Value	5,315	5,889	6,437
Air Quality	267	275	284
Climate Regulation	649	742	806
Erosion Control	103	113	123
Protection from Extreme Events	4,085	4,483	4,881
Food	3,686	4,063	4,446
Pollination	109	120	131
Raw Materials	171	173	177
Recreation	1,258	1,423	1,550
Soil Formation	1,295	1,277	1,279
Waste Treatment	2,934	3,233	3,537
Water Supply	3,088	3,412	3,730
Water Flow Regulation	3,979	4,325	4,682

Oklahoma

Table OK-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Oklahoma

Land use / Land Management Category	Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:	157,666	1,726	157,666	1,785	157,666	1,843
Protected Areas (GAP 1&2)	29,328	533	38,348	697	47,369	862
Protected in Baseline	29,328	533	29,328	533	29,328	533
Added in Scenario	0	0	9,021	165	18,041	329
Converted to Urban Uses	0	0	192	1	176	1
Urban Open Space	0	0	116	0	106	0
Urban Other	0	0	76	0	69	0
SMZs outside protected areas	13,246	241	12,372	225	32,909	598
SMZs w/in protected areas (a)	1,257	(a)	2,121	(a)	9,031	(a)
Available for timber harvest	115,092	953	106,754	862	77,212	383
Ecological Forestry	67	1	2,170	40	1,724	31
Intensive Forestry	66,919	78	60,177	70	56,970	67
Plantation Forestry	0	0	3,167	4	2,998	4
Benign Neglect	48,106	873	41,239	749	15,520	282
Notes:						
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.						

Table OK-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Oklahoma.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	1,726	1,785	1,843
Aesthetic Value	340	353	365
Air Quality	17	18	18
Climate Regulation	41	43	45
Erosion Control	7	7	7
Protection from Extreme Events	262	271	280
Food	236	244	253
Pollination	7	7	7
Raw Materials	11	12	12
Recreation	80	84	87
Soil Formation	84	84	84
Waste Treatment	188	195	202
Water Supply	197	205	212
Water Flow Regulation	255	263	271

South Carolina

Table SC-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for South Carolina

Land use / Land Management Category			Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
			Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:			3,152,309	39,610	3,152,309	42,206	3,152,309	44,684
	Protected Areas (GAP 1&2)		148,875	2,703	636,013	11,588	1,123,152	20,473
		Protected in Baseline	148,875	2,703	148,875	2,703	148,875	2,703
		Added in Scenario	0	0	487,139	8,885	974,277	17,770
Converted to Urban Uses			0	0	55,184	193	50,131	175
		Urban Open Space	0	0	33,392	112	30,334	102
		Urban Other	0	0	21,793	81	19,797	74
SMZs outside protected areas			127,572	2,317	108,973	1,980	320,677	5,823
SMZs w/in protected areas (a)			2,652	(a)	19,749	(a)	130,952	(a)
Available for timber harvest			2,875,863	34,589	2,352,139	28,444	1,658,349	18,213
		Ecological Forestry	40,803	744	75,606	1,379	95,749	1,746
		Intensive Forestry	1,037,899	1,213	798,011	933	665,816	778
		Plantation Forestry	0	0	42,001	49	35,043	41
		Benign Neglect	1,797,161	32,632	1,436,521	26,084	861,741	15,647
Notes:								
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.								

Table SC-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, South Carolina.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	39,610	42,206	44,684
Aesthetic Value	7,913	8,495	9,028
Air Quality	364	372	380
Climate Regulation	966	1,075	1,136
Erosion Control	152	162	172
Protection from Extreme Events	6,036	6,429	6,816
Food	5,488	5,857	6,230
Pollination	162	173	183
Raw Materials	180	189	206
Recreation	1,874	2,059	2,181
Soil Formation	1,689	1,660	1,663
Waste Treatment	4,367	4,659	4,955
Water Supply	4,595	4,918	5,228
Water Flow Regulation	5,824	6,159	6,506

Tennessee

Table TN-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Tennessee

Land use / Land Management Category			Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
			Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:			626,865	7,829	626,865	8,295	626,865	8,754
	Protected Areas (GAP 1&2)		106,375	1,932	189,466	3,447	272,558	4,963
	Protected in Baseline		106,375	1,932	106,375	1,932	106,375	1,932
	Added in Scenario		0	0	83,092	1,516	166,183	3,031
	Converted to Urban Uses		0	0	5,793	20	5,009	18
	Urban Open Space		0	0	3,505	12	3,031	10
	Urban Other		0	0	2,288	8	1,978	7
	SMZs outside protected areas		25,412	462	20,951	381	88,717	1,611
	SMZs w/in protected areas (a)		1,947	(a)	5,895	(a)	52,737	(a)
	Available for timber harvest		495,078	5,436	410,654	4,447	260,580	2,163
	Ecological Forestry		3,574	65	11,674	213	10,934	199
	Intensive Forestry		209,175	244	168,318	197	143,680	168
	Plantation Forestry		0	0	8,859	10	7,562	9
	Benign Neglect		282,329	5,126	221,804	4,027	98,404	1,787
Notes:								
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.								

Table TN-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Tennessee.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	7,829	8,295	8,754
Aesthetic Value	1,563	1,666	1,765
Air Quality	72	74	75
Climate Regulation	191	207	219
Erosion Control	30	32	34
Protection from Extreme Events	1,193	1,264	1,336
Food	1,084	1,152	1,221
Pollination	32	34	36
Raw Materials	36	38	40
Recreation	370	400	422
Soil Formation	336	333	333
Waste Treatment	863	916	971
Water Supply	908	966	1,023
Water Flow Regulation	1,151	1,214	1,278

Texas

Table TX-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Texas

Land use / Land Management Category			Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
			Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:			3,761,857	53,850	3,761,857	56,302	3,761,857	58,730
	Protected Areas (GAP 1&2)		166,676	3,026	924,912	16,856	1,683,147	30,686
	Protected in Baseline		166,676	3,026	166,676	3,026	166,676	3,026
	Added in Scenario		0	0	758,235	13,830	1,516,471	27,660
	Converted to Urban Uses		0	0	16,340	57	14,813	52
	Urban Open Space		0	0	9,887	33	8,963	30
	Urban Other		0	0	6,453	24	5,850	22
	SMZs outside protected areas		274,981	4,995	227,236	4,128	481,039	8,734
	SMZs w/in protected areas (a)		7,142	(a)	53,475	(a)	275,519	(a)
	Available for timber harvest		3,320,200	45,829	2,593,369	35,261	1,582,858	19,258
	Ecological Forestry		35,554	648	94,833	1,730	102,416	1,868
	Intensive Forestry		851,179	995	661,877	774	530,766	620
	Plantation Forestry		0	0	34,836	41	27,935	33
	Benign Neglect		2,433,466	44,186	1,801,824	32,717	921,741	16,737
Notes:								
	a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.							

Table TX-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Texas.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	53,850	56,302	58,730
Aesthetic Value	10,887	11,417	11,935
Air Quality	457	464	473
Climate Regulation	1,329	1,405	1,467
Erosion Control	208	217	227
Protection from Extreme Events	8,245	8,619	8,994
Food	7,552	7,909	8,270
Pollination	222	232	243
Raw Materials	148	189	227
Recreation	2,579	2,718	2,840
Soil Formation	2,016	2,007	2,008
Waste Treatment	6,006	6,290	6,576
Water Supply	6,318	6,621	6,921
Water Flow Regulation	7,883	8,212	8,548

Virginia

Table VA-1. Acreage and Ecosystem Service Value, by Land Use/Land Management Category and Scenario for Virginia

Land use / Land Management Category			Baseline (2016)		Business as Usual (2050)		Conservation (2050)	
			Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)	Acres	Ecosystem Service Value per Year (millions of 2016\$)
Wetland Forest Total:			927,924	13,870	927,924	14,523	927,924	15,131
	Protected Areas (GAP 1&2)		67,211	1,220	266,047	4,847	464,883	8,474
	Protected in Baseline		67,211	1,220	67,211	1,220	67,211	1,220
	Added in Scenario		0	0	198,836	3,627	397,672	7,253
	Converted to Urban Uses		0	0	43,496	152	40,483	142
	Urban Open Space		0	0	26,319	88	24,496	82
	Urban Other		0	0	17,177	64	15,987	59
	SMZs outside protected areas		101,661	1,846	71,742	1,303	117,527	2,134
	SMZs w/in protected areas (a)		1,618	(a)	27,770	(a)	140,337	(a)
	Available for timber harvest		759,052	10,803	546,640	8,221	305,031	4,382
	Ecological Forestry		10,739	196	22,309	407	23,681	432
	Intensive Forestry		175,436	205	95,436	112	64,812	76
	Plantation Forestry		0	0	5,023	6	3,411	4
	Benign Neglect		572,878	10,402	423,871	7,696	213,127	3,870
Notes:								
a. Acreage and ecosystem service value in SMZ's that are also within protected areas are included in the subtotals for Protected Areas. Acreage is provided here for reference.								

Table VA-2: Ecosystem Service Value of Wetland Forests, by Ecosystem Service and Scenario, Virginia.

Ecosystem Service	Baseline (2016)	Business as Usual (2050)	Conservation (2050)
	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)	Ecosystem Service Value per Year (millions of 2016\$)
Total:	13,870	14,523	15,131
Aesthetic Value	2,814	2,971	3,100
Air Quality	115	116	118
Climate Regulation	344	392	406
Erosion Control	54	56	58
Protection from Extreme Events	2,127	2,218	2,313
Food	1,952	2,033	2,125
Pollination	57	60	62
Raw Materials	31	35	46
Recreation	667	741	769
Soil Formation	497	474	476
Waste Treatment	1,552	1,617	1,689
Water Supply	1,633	1,713	1,788
Water Flow Regulation	2,027	2,096	2,182

Appendix 2: Candidate Ecosystem Service Productivity Values for Land-Use and Ecosystem Service Combinations

As explained under [Methods Ecosystem Service Valuation](#), the benefit transfer method applies estimates of ecosystem service value from existing studies of “source areas” to the “policy area,” which in this case comprises wetland forests in the the various land use/land management classes identified through spatial and other analysis. This application is done on a land-use-by-land-use basis. The table below lists all of the values from source area studies considered for our calculations. All values have been adjusted for inflation to reflect 2016 price levels. For source area values originally reported in currencies other than U.S. dollars, the values have been converted using exchange rates.

Biome	Ecosystem Service	Minimum 2016\$ per acre per year	Maximum 2016\$ per acre per year	Source
Forest	Aesthetic	62.66	850.48	Moore, R. (2011)
		4,501.06	18,392.66	Nowak, D J, Crane, D E, Dwyer, J F. (2002)
	Air Quality	227.48	227.48	Weber, Ted (2007)
	BioControl	2.57	2.57	Brenner-Guillermo, J. (2007)
	Climate	68.39	68.39	Brenner-Guillermo, J. (2007)
		57.67	57.67	Costanza, R, R d’Arge, S Farber, M Grasso, R deGroot, B Hannon, and M van den Belt. (1997)
		3.32	46.21	Earth Economics Database
		4.37	4.37	Hall, D.S. (2010)
		328.24	328.24	Flores, L., Harrison-Cox, J., Wilson, S., & Batker, D. (2013)
	Erosion	62.73	62.73	Brenner-Guillermo, J. (2007)
		3.14	36.59	Zhou, X, Al-Kaisi, M, Helmers, J M. (2009)
	Extreme Events	316.59	316.59	Moore, R. (2011)
		808.68	808.68	Weber, Ted (2007)
	Food	2,453.84	2,453.84	Kauffman, G., Homsey, A., McVey, E., Mack, S., and Chatterson, S. (2011)
	Pollination	205.67	205.67	Brenner-Guillermo, J. (2007)
		1.43	7.20	Costanza, R. Wilson, M.Troy, A.Voinov, A.Liu, S.D’Agostino, J. (2007)
		89.32	89.32	Weber, Ted (2007)
	Raw Materials	21.31	21.31	Hall, D.S. (2010)
		169.12	169.12	Weber, Ted (2007)
	Recreation	154.77	154.77	Brenner-Guillermo, J. (2007)
		1.31	4.62	De la Cruz, A. and J. Benedicto (2009)
		4.00	4.00	Haener, M K, Adamowicz, Wiktor L. (2000)
		3.78	3.78	Hall, D.S. (2010)
		37.64	46.13	Prince, R, Ahmed, E. (1989)
		2.82	510.93	Shafer, E L, Carline, R, Guldin, R W, Cordell, H K.

				(1993)
	Soil fertility	6.17	6.17	Brenner-Guillermo, J. (2007)
		20.25	20.25	Weber, Ted (2007)
	Waste	56.04	56.04	Brenner-Guillermo, J. (2007)
		8.78	8.78	De la Cruz, A. and J. Benedicto (2009)
		269.46	270.58	Zhongwei, Lui. (2006)
	Water	207.21	207.21	Brenner-Guillermo, J. (2007)
		48.04	48.04	De la Cruz, A. and J. Benedicto (2009)
		11.08	11.08	Hall, D.S. (2010)
		316.59	316.59	Moore, R. (2011)
		1,310.08	1,310.08	Weber, Ted (2007)
	Water Flows	233.18	233.18	Mates, William. (2007)
		808.68	808.68	Weber, Ted (2007)
Urban Open Space	Aesthetic	461.00	461.00	Johnston, Robert J, Mazzotta, Marisa J, Opaluch, James J, Grigalunas, Thomas A. (2001)
		1,751.15	3,027.76	Opaluch, J, Grigalunas, T, Mazzotta, M, Johnston, R, Diamantedes, J. (1999)
		1,019.96	1,340.58	Qiu, Z, Prato, T, Boehm, G. (2006)
	Air Quality	32.91	32.91	McPherson, E G, Scott, K I, Simpson, J R. (1998)
		195.01	195.01	McPherson, G. (1992)
	Climate	1,150.06	1,150.06	McPherson, G. (1992)
	Extreme events	319.88	605.26	Streiner, C, Loomis, J. (1995)
	Water Flows	8.44	8.44	McPherson, G. (1992)
		140.13	190.17	Trust for Public Land. (2010)
Urban Other	Climate	426.76	426.76	Brenner-Guillermo, J. (2007)
		69.81	69.81	Moore, R. (2011)
	Recreation	2,707.64	2,707.64	Brenner-Guillermo, J. (2007)
	Water	1.28	1,501.62	Moore, R. (2011)
	Water flows	7.71	7.71	Brenner-Guillermo, J. (2007)
Wetland	Aesthetic	38.99	38.99	Amacher, G.S., R.J. Brazee, J.W. Bulkley and R.A. Moll (1989)
		1,829.89	1,829.89	Gerrans, P. (1994)
		972.18	1,557.57	Johnston, Robert J, Mazzotta, Marisa J, Opaluch, James J, Grigalunas, Thomas A. (2001)
		5,993.29	14,207.18	Mazzotta, M. (1996)
		9.89	117.62	Moore, R. (2011)
		7,078.85	9,549.09	Opaluch, J, Grigalunas, T, Mazzotta, M, Johnston, R, Diamantedes, J. (1999)
		468.10	1,497.91	Thibodeau, Francis R, Ostro, Bart D. (1981)
	Air Quality	76.54	99.37	Jenkins, W A, Murray, B C, Kramer, R A, Faulkner, S P. (2010)
	Climate	159.91	159.91	Brenner-Guillermo, J. (2007)
		0.89	14.47	EarthEconomics Database

		218.26	218.26	Hughes, Zachary. (2006)
		80.12	584.73	Jenkins, A.W. (2009)
		637.40	637.40	Moore, R. (2011)
		63.55	4,929.88	Wilson, Sara. (2013)
	Erosion	179.84	179.84	Weber, Ted (2007)
	Extreme events	111.58	4,646.59	Brenner-Guillermo, J. (2007)
		308.38	308.38	Costanza, R, Farber, S C, Maxwell, J. (1989)
		282.63	282.63	Costanza, R, Farley J. (2007)
		2,189.84	2,189.84	Costanza, R., R. d'Arge, R. de Groot, S. Farber, M. Grasso, B. Hannon, K. Limburg, S. Naeem, R.V. O'Neill, J. Paruel, R.G. Raskin, P. Sutton and M. van den Belt (1997)
		8,543.79	8,543.79	Gupta, T.R. and J.H. Foster (1975)
		1,668.33	7,617.80	Leschine, Thomas M, Wellman, Katherine F, Green, Thomas H. (1997)
		864.22	864.22	Moore, R. (2011)
		6,241.32	6,241.32	Thibodeau, Francis R, Ostro, Bart D. (1981)
	Food	17,113.98	17,113.98	Aburto-Oropeza, O, Ezcurra, E, Danemann, G, Valdez, V, Murray, J, Sala, E. (2008)
		5.82	729.18	Batie, Sandra S, Wilson, James R. (1978)
		60.71	141.86	Bell, Frederick W. (1989)
		194.93	194.93	Gren, I.M. and T. Soderqvist (1994)
		3,371.29	3,371.29	Hamilton, L.S. and S.C. Snedaker (1984)
		826.69	826.69	Hughes, Zachary. (2006)
		1,030.72	1,030.72	Johnston, Robert J, Mazzotta, Marisa J, Opaluch, James J, Grigalunas, Thomas A. (2001)
	Raw materials	50.86	50.86	Everard, M. (2009)
		86.38	86.38	Kosz, M., B. Brezina and T. Madreiter (1992)
	Recreation	429.95	429.95	Anderson, Glen D, Edwards, Steven F. (1986)
		157.38	1,207.49	Bell, Frederick W. (1989)
		81.83	81.83	Bergstrom, John C, Stoll, John R, Titre, John P, Wright., Vernon L. (1990)
		1,740.48	1,786.24	Brenner-Guillermo, J. (2007)
		110.81	435.91	Costanza, R, Farber, S C, Maxwell, J. (1989)
		1,055.42	1,055.42	Creel, Michael, Loomis, John. (1992)
		89.28	1,008.24	Gren, I.M. and T. Soderqvist (1994)
		221.81	1,699.20	Jaworski, E, Raphael, C N. (1978)
		7.50	7.50	Jenkins, A.W. (2009)
		210.88	210.88	Kreutzwiser, R. (1981)
		212.40	212.40	Lant, C.L. and R.S. Roberts (1990)
		585.87	10,504.13	Thibodeau, Francis R, Ostro, Bart D. (1981)
		124.40	124.40	Whitehead, J C, Groothuis, P A, Southwick, R, Foster-Turley, P. (2009)
		657.53	4,261.90	Whitehead, J C. (1990)

	Soil fertility	2,149.72	2,149.72	Bystrom, O. (2000)
	Soil Formation	535.94	535.94	Weber, Ted (2007)
	Waste	67.95	6,398.80	Breaux, A, Farber, S, Day, J. (1995)
		1,064.85	1,064.85	Brenner-Guillermo, J. (2007)
		172.40	172.40	Gren, I.M. and T. Soderqvist (1994)
		558.63	558.63	Jenkins, W A, Murray, B C, Kramer, R A, Faulkner, S P. (2010)
		212.40	212.40	Lant, C.L. and R.S. Roberts (1990)
		4,620.66	4,620.66	Thibodeau, Francis R, Ostro, Bart D. (1981)
		11,031.50	11,031.50	Zhongwei, Lui. (2006)
	Water	1,961.57	2,440.78	Brenner-Guillermo, J. (2007)
		631.37	631.37	Creel, M, Loomis, J. (1992)
		18.44	18.44	Folke (1991)
		211.98	849.20	Moore, R. (2011)
		18,861.26	18,861.26	Thibodeau, Francis R, Ostro, Bart D. (1981)
		1,310.08	1,310.08	Weber, Ted (2007)
	Water flows	3,793.57	3,793.57	Brenner-Guillermo, J. (2007)
		3,974.86	3,974.86	Leschine, T.M., K.F. Wellman and T,H. Green (1997)
		4,389.53	4,389.53	UK Environment Agency (1999)